ISSN 1447-4999

Australian Prehospital Emergency Health Research Forum

Peer-Reviewed

ABSTRACTS

from

The Australian College of Ambulance Professionals (ACAP)

2010 Conference

Perth, Western Australia

14-16 October 2010

The Journal of Emergency Primary Health Care Management Committee gratefully acknowledges the support of ACAP, and all authors who submitted abstracts for peer review to the Australian Prehospital Emergency Health Research Forum (APEHRF) and further presentation at the ACAP 2010 Conference in Western Australia.

Additionally, the Management Committee sincerely thanks the following peer reviewers and adjudicators for volunteering their valuable time and expertise in the peer review of abstracts, evaluation of posters or adjudication of selected oral presentations at the Conference, from which their collective results determined the winners of the 2010 APEHRF (David Komesaroff) Best Paper Award; APEHRF Best Paper in the category of Higher Degree by Research; APEHRF Best Paper in the category of Undergraduate Student; and APEHRF Best Poster Award.

Award Winners:

APEHRF (David Komesaroff) Best Paper Award

Kate Cantwell, Janet Bray, Michael Stephenson, Kerry Power, Karen Smith (VIC)
Importance of pre-hospital blood pressure post ROSC on survival to hospital discharge.

Best Paper in the category of Higher Degree by Research

Brett Williams, Andrys Onsman, Ted Brown (VIC)
Validation of the paramedic graduate attribute scale (PGAS): a RASCH rating analysis.

Best Undergraduate Student Paramedic Paper

Pauline Murcott, Brett Williams, Amee Morgans, Malcolm Boyle (VIC)
Community perceptions of the professional status of the paramedic discipline.

Best Poster Prize

Jennifer Melvin, Brett Williams Malcolm Boyle (WA, VIC)
Pre-shock CPR: should we do it?

Abstract Peer Reviewers:

Mr. Stephen Burgess (VIC), Mr. Joseph Cuthbertson (WA), Prof. Gerry FitzGerald (QLD), Mr. Paul Jennings (VIC), Prof. Peter O'Meara (NSW), Prof. Helen Snooks (UK), A/Prof. Vivienne Tippet (QLD).

Poster Adjudicators:

Mr. Chris Cotton (SA), Mr. Alan Eade (VIC), Mr. John Hall (NSW), Mr. Chris Huggins (VIC), Dr. Harry Oxer (WA), Mr. Doug Wright (ACT).

Conference Adjudicators:

Mr. Joe Acker (NSW) Mr. Jeff Allan (VIC), Dr. Jason Bendall (NSW), Mr. Chris Cotton (SA), Mr. Alan Eade (VIC), Dr. Cindy Hein (SA), Mr. Chris Huggins (VIC), Mr. Paul Jennings (VIC), Mr. Mick Lazell (QLD), Ms. Tammy Lee (TAS), Mr. Bill Lord (VIC), Dr. Paula McMullen (TAS), Prof. Peter O'Meara (NSW), Dr. Harry Oxer (WA), Dr. Brian Sengstock (QLD), Mr. Doug Wright (ACT).

PREVENTING THE TRANSMISSION OF INFECTIONS IN THE PARAMEDIC WORK SETTING: ARE THE CURRENT NATIONAL INFECTION CONTROL GUIDELINES ADEQUATE?

Nigel Barr BN, Grad Dip Advanced Clinical Nursing (Intensive Care), AsDip HlthSc(Ambulance Officer), AdDip HlthSc(MICA Studies) **Mark A Holmes** BSc(Hons), PhD **Anne Roiko** BSc(Hons), PhD **Nicolas J Prass** BHLthSc, MHLthSc, Grad Dip Intensive Care Paramedic

University of the Sunshine Coast, School of Health and Sport Sciences, Sippy Downs Qld
Australia

Background

Considerable research has been conducted into appropriate infection control practices for health-care establishments such as hospitals and long-term care facilities. Infection control practices for these health-care settings have been well described in national guidelines, such as the one published by the Department of Health and Aging (DoHA). The paramedic work setting, however, can present unique challenges for infection control as it involves invasive clinical procedures that are often performed in a mobile or uncontrolled environment. While paramedics apply best practice with infection control to ensure personal and patient safety, there has been a paucity of research into the potential for transmission of pathogens in the paramedic work setting.

Objective

To evaluate the adequacy of current infection control guidelines for their application in the paramedic work setting.

Methods

A comprehensive desktop analysis of the infection control guidelines published by DoHA and those produced and supplied by Australian and New Zealand ambulance authorities was performed.

Findings

The analysis of the DoHA infection control guidelines found little recognition of the unique hazards present in the paramedic work setting. In contrast, the guidelines produced by the ambulance authorities were focussed more on managing the risk of infection transmission for paramedics, but there were some differences in the infection control practices being recommended. It is clear from the analysis that the issues requiring further consideration to minimise the risk of transmission of pathogens in the paramedic work setting include: 1) vehicle design and cleaning; 2) use of portable equipment in an uncontrolled environment; 3) treatment of multiple patients with limited infection control resources; and 4) governance of specialised procedures and surveillance programs for the pre-hospital emergency environment.

Conclusions

This study has highlighted the need for further research to determine whether standard precautions for infection control developed for health-care establishments are also effective in the paramedic work setting. Major risk factors need to be determined that are specific to paramedic practice so that appropriate quality management procedures for infection control can be developed.

THE PREHOSPITAL MANAGEMENT OF CROUP: A LITERATURE REVIEW

Chloe Bell (BMus) **Christian Winship, Erin Sorensen, Brett Williams** MHLthSc, PhD
Candidate **Malcolm Boyle** MCLinEpi, PhD

Department of Community Emergency Health and Paramedic Practice, Monash University

Background

Croup or acute laryngotracheobronchitis is an acute upper respiratory syndrome that is viral in origin, and is most prevalent in children between the ages of six months and five years. The major concern for paramedics is when croup presents as full or partial obstruction of the upper airway due to subglottic oedema.

Objectives

The objective of this study was to identify the best practice management for croup in the prehospital setting.

Methods

A literature review was undertaken of the electronic medical databases Ovid Medline, EMBASE, CINAHL Plus, Cochrane Systematic Review, and Meditext. Keywords used in the search included; croup, acute laryngotracheobronchitis, EMS, EMT, paramedic, emergency medical service, emergency medical technician, prehospital, and ambulance. The keywords were used individually and in combination. Articles were included if they reported on prehospital management of croup or acute laryngotracheobronchitis. Articles not written in English were excluded.

Findings

Of the 25 articles located, only three low level evidence articles met the inclusion criteria. References of relevant article were also reviewed with no additional articles identified. Of the three articles, one case study highlighted the effective use of heliox in the treatment of croup by an air ambulance service. The other two case studies contained only brief comments on the use of nebulised adrenaline, or nebulised budesonide, or oral dexamethasone to treat croup in the prehospital setting. From non-prehospital articles identified in the search the prehospital management of croup appears to be based on findings from hospital-based studies.

Conclusion

There is a lack of evidence supporting the current prehospital management of croup, with current management based on hospital studies. There is a need for clinical research to identify an evidence-based approach to croup management in the prehospital setting.

EPIDEMIOLOGY OF ENDOTRACHEAL INTUBATION IN A LARGE AUSTRALIAN AMBULANCE JURISDICTION

Jason Bendall, Jillian Patterson and Paul Middleton

Ambulance Research Institute, Ambulance Service of New South Wales

Background

There is longstanding controversy surrounding paramedic performed out-of-hospital intubation, and little is known about the epidemiology of non drug-facilitated intubation attempts within the Ambulance Service of NSW.

Objective

To describe the epidemiology of out-of-hospital intubation attempts in a large Australian ambulance service.

Methods

All unique cases were identified where an intubation attempt was documented on a patient health care record by intensive care paramedics, between 1 July 2007 and 30 June 2008.

Findings

There were 1577 unique cases identified, and intubation attempts were documented as successful in 87% of cases. The median number of intubation attempts during this period was 1 (IQR 0-3) and the majority of intubation attempts were for cardiac arrests (79%). Intubated patients were aged 10 or over in 98% of all attempts; however the majority of ICP (88%) did not attempt intubation in a patient < 16 years. There was no evidence that success rates in patients aged < 10 were lower than patients aged ≥ 10 ($p=0.55$). Only 25 operational intensive care paramedics (<3%) attempted 10 or more intubations during the study period, and approximately 40% of ICP did not attempt intubation in the year of study.

Conclusion

Intubation is an uncommonly performed intensive care paramedic procedure, the majority of attempts are for cardiac arrest and the majority of attempts are successful, however very few intensive care paramedics perform 10 or more intubations annually. Success rates do not appear to differ between paediatric and adult populations despite very few intubation attempts occurring in children.

EPIDEMIOLOGY OF INCIDENTS INVOLVING OLDER PATIENTS WHO HAVE FALLEN AND CALLED FOR AN EMERGENCY AMBULANCE RESPONSE

Jason Bendall, Jillian Patterson, Paul Simpson, Paul Middleton

Ambulance Research Institute, Ambulance Service of New South Wales

Background

Provision of emergency care to an aging population is a core function of modern ambulance jurisdictions. Falls in this population are one of the most common reasons for an older patient to call triple zero. Despite the high incidence of falls-related cases, little is known about the nature of these incidents. Describing older fallers attended to by ambulance will likely inform policy and practice that will optimise ambulance management of this vulnerable population.

Objective

To describe the epidemiology of older fallers attended to by paramedics.

Methods

A retrospective analysis was conducted of computer-aided dispatch (CAD) and patient health care record (PHCR) databases for all calls assigned a medical priority dispatch system (MPDS) category of “Falls” in the period 1 July 2008 to 30 June 2009 for patients aged 65 years or more.

Findings

Overall there were 73,550 calls classified as ‘Falls’, of which 45,117 (61%) were to patients aged 65 years or more. The age standardized rate for falls in this cohort of patients was 4311.2 per 100,000 population (95% CI 4271.2-4351.5). Falls most commonly occurred mid-morning between 0900 and 1200, and peaked on Fridays. The vast majority of calls occurred in metropolitan areas (67.8%) compared to regional/remote settings. Falls resulted in injury in 53% of cases, with ‘limb injuries and fractures’ (59%), ‘head injury’ (19%) being the most common trauma protocol recorded. The non-transport rate for older fallers was 25%.

Conclusions

Falls in patients aged 65 years or more represent a huge clinical and operational burden for modern ambulance jurisdictions. Further prospective research is required to accurately identify and describe this at-risk population of patients.

PARAMEDIC PERCEPTIONS OF REASONS FOR PATIENTS TO BE CONVEYED TO AN EMERGENCY DEPARTMENT

**Jason Bendall, Paul Simpson, Monika Sitkowski, Suzanne Davies,
Jillian Patterson, Paul Middleton**

Ambulance Research Institute, Ambulance Service of New South Wales

Background

All ambulance jurisdictions are facing operational pressures, including increasing demand, aging populations and emergency department ramping. Little is known about paramedic perceptions of ambulance utilisation for patients conveyed to an emergency department (ED).

Objectives

To describe paramedic perceptions of reasons for patients to be conveyed by ambulance to an emergency department.

Methods

A survey of paramedics was undertaken over a 24 hour period in January 2010. Paramedics completed a 12 statement questionnaire using Likert scales to explore perceptions of the factors influencing disposition. Paramedic responses and patient health care records were analysed using SAS v 9.1.3.

Findings

During the 24-hour study period 1518 patients were transported to an emergency department, and there were 850 surveys returned (56.0%). Paramedics agreed that patients required acute (urgent) ED care in 37% of cases, and less urgent ED care in 46% of cases. Paramedics agreed that patients could have their needs met by another health service in 38% of cases. Factors that paramedics felt influenced ED disposition were - patient / carer concern (69%); patient requesting transport (57%); patient unable to see GP (35%); and patient not having own transport (36%). Paramedics agreed that patient not having a GP influenced transport in only 11% of cases.

Conclusions

Paramedics agree that the majority of patients transported need ED care however there is a perceived opportunity to explore alternatives to ED care for some patients. Patient or carer concern and patients requesting transport appear to be common in transported patients. Further study is required to validate paramedic perceptions.

MULTIDISCIPLINARY RESPONSES TO MENTAL HEALTH CRISES IN THE PRE-HOSPITAL SETTING

Leanne Boyd, Jade Sheen, Matthew Johnson, Kathryn Eastwood MEmerHlth (MICA),
GradDipEmerHlth (MICA), BSc, RN, BParamedicStudies, DipAmbParamedStud

Department of Community Emergency Health and Paramedic Practice, Monash University,
Melbourne

Background

Consumer rights, suicide management and continuity of care have been raised as national mental health priorities. A lack of teamwork and interagency liaison, limited adherence to the Mental Health Act (MHA), limited de-escalation skills and a limited understanding of the experiences of consumers are systemic issues recognised within the recent Victorian Auditor-General report.

Objectives

This project aimed to address these issues by creating an interactive education package focusing on the multidisciplinary management of mental health crises such as suicide, to be integrated into existing mental health and interprofessional units within the Faculty of Medicine, Nursing and Health Sciences at Monash University.

Methods

An education program with four modules was developed after extensive consultation with key stakeholders. Each module consists of a one hour lecture with embedded video clips and scenarios and a one hour interactive tutorial with activities such as role play, case based learning and group discussion. The program was piloted with Double Degree (Nursing/Paramedic) students. Feedback was obtained through focus groups and surveys.

Findings: Focus group and survey analysis revealed that the program was well received by participants. Strategies to improve the program were identified and modifications were made prior to official implementation of the program.

Conclusions

The preliminary findings show that the program can be successfully integrated into existing curricula of undergraduate health science courses. With continued broader dissemination, this program has the potential to improve outcomes for consumers and their carers during a mental health crisis within the pre-hospital setting. An enhanced understanding of key stakeholder roles and responsibilities has the potential to enhance teamwork, facilitate interagency collaboration and optimize the care of both consumers and their families.

Acknowledgements

The research team would like to acknowledge the assistance of the funding body, Australian Government Department of Health and Ageing.

NATIONALISATION OF THE AMBULANCES FOR ASTHMA PROGRAM

Leanne Boyd DipAppSci, BNursing, GradDipCritCare, MNursing, GradCertHigherEd, PhD
Matthew Johnson DipAmbParamedStudies, GradDipEmergHealth (MICA)

Department of Community Emergency Health and Paramedic Practice, Monash University,
Melbourne

Background

Previous studies revealed that utilisation of ambulance services for asthma is suboptimal, despite improved outcomes when paramedic intervention was sought. The 'Ambulances for Asthma' program is an interactive web-based decision aid to assist people with asthma to recognise when they would benefit from paramedic intervention rather than self transporting to hospital. Initially, the program aimed to appropriately improve ambulance utilisation and knowledge of resources in rural Victoria. Based on positive evaluation outcomes the program received further funding from The Asthma Foundations of Australia to nationalise all content.

Objectives

1. Adapt the existing 'Ambulances for Asthma' program to incorporate relevant information on a state by state basis after extensive consultation with all key stakeholders
2. Increase awareness of the appropriate use of emergency ambulance services for asthma amongst patients and their carers in each Australian state and territory.

Methods

The funding body did not require the research team to formally re-evaluate the 'Ambulances for Asthma' web site as a component of this project. Despite this, a number of measures such as organisational/consumer consultation and human-computer interface testing were undertaken to ensure the quality and effectiveness of the intervention redevelopment. Tracking measures have been implemented to monitor website utilisation for future evaluations.

Findings

The program was given final approval by all key stakeholders prior to going live in 2009.

Conclusions

People with asthma Australia wide can now access asthma information and resources relevant to their state or territory. Vital links to other resources such as the Royal Flying Doctor Service, individual state/territory Ambulance Services and Asthma Foundations have been provided. The website includes relevant information for people with asthma living in metropolitan, rural and remote regions of Australia. A number of additional resources have been included to facilitate people with asthma wishing to travel to remote areas of Australia.

NANNA DOWN AND OTHER FALLS

Malcolm Boyle ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClintEpi, PhD

Monash University, Department of Community Emergency Health and Paramedic Practice,
Melbourne, Australia

Background

The number of elderly patients admitted to hospital emergency departments has increased by approximately 10% per year for the past few years with this number expected to increase with the aging population. The cause of these falls varies from a simple loss of balance to more significant medical conditions. Falls are not just confined to the elderly, trauma from falls is second behind transport related accidents for adults and children. To date no state-wide study has been conducted that has looked at falls and the outcomes from a prehospital perspective.

Objectives

The objective of this study was to identify fall incidents and their outcomes from a prehospital perspective for a 12 month period.

Methods

Fall incidents were identified in the Victorian Prehospital Trauma Triage Study data repository. The fall data was then linked to the Victorian State Trauma Registry to determine patient outcomes.

Findings

There were 24,059 falls identified for the 12 month period, this amounts to approximately 45% of the trauma workload and 6% of the total workload by Ambulance Victoria. Standing falls with no prehospital potential major trauma criteria accounted for 38.7% of the total trauma incidents with 6.7% of falls having one or more of the prehospital potential major trauma criteria. For 2.4% of fall patients who experienced hospital defined major trauma 25% of this group did not have prehospital potential major trauma criteria. 33% of fall patients who did not have prehospital potential major trauma criteria went on to die.

Conclusions

This study demonstrates that falls are a large percentage of the trauma workload, however, a small percentage of falls have no prehospital potential major criteria but go on to have hospital defined major trauma. Further research is required to better identify those patients who fall and have hospital major trauma and not prehospital potential major trauma criteria.

DO MANNEQUIN CHESTS PROVIDE AN ACCURATE REPRESENTATION OF A HUMAN CHEST FOR SIMULATED DECOMPRESSION OF TENSION PNEUMOTHRACES?

Mal Boyle ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClinEpi, PhD
Brett Williams Intensive Care Paramedic Cert, MICA Cert, BAVEd, GCert ICP, GDip EmergHlth, MHIthSc

Monash University, Department of Community Emergency Health and Paramedic Practice,
Melbourne, Australia

Background:

The presentation of traumatic tension pneumothorax in the Victorian prehospital setting is <1%. Therefore, it is important this uncommon presentation, managed by needle decompression, is practised by paramedics and undergraduate paramedics using a range of educationally sound and realistic low, medium, and high fidelity mannequins.

Objectives:

To identify if mannequin chests are an accurate representation of a human chest for simulated decompression of tension pneumothoraces.

Methods:

This is a two part study. Part 1, was a review of the literature to identify chest wall thickness in humans, and Part 2, measurement of the chest wall thickness for two commonly used mannequins. For Part 1, the literature search was conducted using the Cochrane Central Register of Controlled Trials (CENTRAL), MEDLINE, CINAHL, and EMBASE databases from their beginning until the end of March 2010. Keywords included chest wall thickness, tension pneumothorax, pneumothorax, thoracostomy, needle thoracostomy, decompression, and needle test. Studies were included if they reported chest wall thickness. For Part 2, a set of measurement callipers was used to determine the chest wall thickness in the area of the second intercostal space mid clavicular line.

Findings:

Part 1, the literature review located 3,684 articles with 7 meeting the inclusion criteria. Chest wall thickness in adults varied between 3cm and 9.3cm at the area of the second intercostal space mid clavicular line. Part 2, in the area of the second intercostal space mid clavicular line, right side of the chest was 1.1cm thick and 1.5cm on the left for the Laerdal mannequin. The MPL mannequin in the same area, right side of the chest was 1.4cm thick and 1.0cm on the left.

Conclusions:

Mannequin chests are not an accurate representation of the human chest when used for simulated decompression of tension pneumothoraces and therefore may not provide a realistic educational or training experience.

RURAL CLINICAL PLACEMENTS, ARE THEY WORTH IT?

Malcolm Boyle¹ ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClInEpi, PhD
Brett Williams¹ Intensive Care Paramedic Cert, MICA Cert, BAVEd, GCert ICP, GDip
EmergHlth, MHIthSc
Nathan Stam¹ BSc(Hons), DAPS

¹Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

Bachelor of Emergency Health (Paramedic) (BEH) students at Monash University undertake rural clinical placements as part of their on-road placements to assist with the transition from student to novice paramedic. Anecdotally, students report a lack of work, lack of interest shown by the paramedics, and a lack of opportunity to practise their clinical skills whilst on rural placements. These issues have not previously been documented in Australian paramedic literature.

Objectives

The objective of this study was to investigate student's experiences during rural clinical placements.

Methods

A cross-sectional study using a paper-based questionnaire with a convenience sample of second and third year BEH undergraduate students. Ethics approval was granted.

Findings

Eighty four second and third year BEH students participated. 59.5% were female and 75% < 25 years of age. 68% of students identified a difference in the workload mix (incidents attended) between metropolitan and rural placements. 75% of students stated there was > 4 hours downtime in rural compared to 12% for metropolitan placements ($p < 0.0001$). 37% of students stated that crews were willing to participate in downtime activities, i.e. skills practise. Due to the excessive downtime 80% of students thought additional online learning activities would benefit their consolidation of theory into practice. Ambulance crews were aware of the student's role during the placement 19% of the time with 73% of students stating they were not given the opportunity to manage the patient to their level of education.

Conclusions

This study demonstrates that there is more downtime during rural clinical placements, however, this time is poorly utilised for theory practice consolidation. Further effort and collaboration is required by universities and industry to maximise students' learning during clinical placements.

RAPID SEQUENCE INTUBATION BY PARAMEDICS: IMPROVING CARE WITH PROPOFOL

Nicholas Bridge¹ Michael Adam² Dr. John Maloney³

¹Intensive Care Flight Paramedic - Air Ambulance Victoria ²Intensive Care Flight Paramedic Team Manager - Air Ambulance Victoria ³Consultant Anaesthetist, Alfred Hospital, Melbourne.

Background

Intensive Care Flight Paramedics (MFP) at Air Ambulance Victoria (AAV) commonly practice Rapid Sequence Intubation (RSI) using Fentanyl, Midazolam and Suxamethonium. Literature review demonstrates limited pre-hospital use of Propofol, if any, by Paramedics for RSI. The AAV prescribed adult doses of Fentanyl and Midazolam for RSI in conscious (GCS>12) burns patients on occasions fails to completely sedate patients prior to neuromuscular blockade administration. In 2008 Propofol was authorised for use by AAV MFP in RSI.

Objectives

To identify frequency, effectiveness and complications associated with Propofol use in pre-hospital RSI by MFP in conscious, airway burn injury.

Methods

Data was gathered from patient care records from primary response helicopters across Victoria from January 2008.

Inclusion criteria: Actual or potential airway compromise from burns requiring intubation by MFP. Adults with GCS >12.

Exclusion criteria: Injuries in addition to or other than burns.

Analysis included physiological parameters pre and post intubation, requirement for additional pre-intubation sedation after the initial induction dose, airway grade view and intubation success.

Findings

In 17 cases Propofol (1-1.5mg/kg) was utilised for induction in RSI. In 3 cases a further 0.5mg/kg was required to achieve sedation. There were no significant differences between pre and post induction HR or BP ($p=0.59$ and 0.40 respectively). Intubation was achieved on first attempt in 100% of cases with grade 2 or better view.

Conclusions

Intensive Care Flight Paramedic use of Propofol in RSI, enhances clinical care and provides effective sedation prior to intubation in conscious burns patients, without adverse haemodynamic consequences.

HANGING ASSOCIATED OUT-OF-HOSPITAL CARDIAC ARREST; A REVIEW OF 1,266 EMS CASES FROM THE VICTORIAN AMBULANCE CARDIAC ARREST REGISTRY (VACAR)

**Dr Conor Deasy, MB, BAO, BCH, FCEM¹⁻³ Janet Bray, MCH² Dr Karen Smith, PhD
Dr Linton Harris, PhD A/Prof Steve Bernard, MD, FJFICM¹⁻³ Prof Peter Cameron, MD,
FACEM^{1,3}**

on behalf of the **VACAR Steering Committee.**

¹Monash University Department of Epidemiology and Preventive Medicine ²Ambulance
Victoria ³Alfred Hospital

Introduction

Hanging is an infrequent but devastating cause of out-of-hospital cardiac arrest (OHCA). We sought to determine the characteristics and outcomes of hanging associated OHCA in Melbourne Australia.

Methods

A 10-year retrospective case review of all adult hangings (aged ≥ 16 years) associated with OHCA, was performed using data from the Victorian Ambulance Cardiac Arrest Registry (VACAR).

Results

Between 2000 and 2009 EMS attended 1266 cases of OHCA where hanging was the precipitant. These cases represented 4.1% (range 3.0% to 4.9%) of all OHCA annually. Patients were mostly male (78%) and the mean age was 41 years (range 16-99 years). Overall bystander CPR was performed in 14% of cases (n=179) and in 48% of cases (n=112 of 235) where EMS attempted resuscitation. The most common initial rhythm was asystole (96.5%), followed by PEA (3%) and VF (0.5%). Where EMS attempted resuscitation (n=235), a sustained return of spontaneous circulation (ROSC) at scene was achieved in 64 (27%) patients, 7 (3%) surviving to hospital discharge. Of these 7 survivors, all occurred after 2006, 3 (43%) had bystander CPR, and two were in VF (29%).

Discussion

ROSC rates are low in this group of patients and overall survival is rare. Nearly one third of survivors had an initial rhythm of VF which supports early application of defibrillator pads to ascertain a presence of a shockable rhythm. Given the poor survival rate, pre-hospital guidelines and training should highlight the social and pastoral needs of the victim's families and witnesses to these distressing cases.

OUT-OF-HOSPITAL CARDIAC ARRESTS IN OCTOGENARIANS AND NONAGENARIANS IN MELBOURNE

**Dr Conor Deasy, MB, BAO, BCH, FCEM¹⁻³ Janet Bray, MCH² Dr Karen Smith, PhD
Dr Linton Harris, PhD A/Prof Steve Bernard, MD, FJFICM¹⁻³ Prof Peter Cameron, MD,
FACEM^{1,3}**

¹Monash University Department of Epidemiology and Preventive Medicine ²Ambulance
Victoria ³Alfred Hospital

Introduction

Controversy exists around CPR in the elderly. We examine the characteristics and outcomes of out-of-hospital cardiac arrest (OHCA) in elderly patients in Melbourne, Australia.

Methods

The Victorian Ambulance Cardiac Arrest Registry (VACAR) was searched for all OHCA's occurring in patients aged 80-89 and 90-99 years, that were unwitnessed by EMS.

Results

Between 2000 and 2009 there were 30,008 OHCA's of which 6,430 (21.4%) and 1,530 (5.1%) were in octogenarians and nonagenarians respectively. The most common cardiac rhythm on arrival of EMS for both octogenarians and nonagenarians was asystole (74.6% and 79.3%, respectively), PEA (14.2% and 14.8%, respectively) and VF/VT in (11.2% and 5.9%, respectively). Resuscitation was attempted in 39.5% (ten year range 37.7% - 42.4%) of octogenarians and 31.6% (ten year range 23.5% - 39.1%) of nonagenarians. Where EMS attempted resuscitation ROSC rates have increased from 19% to 34% ($p < 0.001$) but was not significantly different for nonagenarians (16%, $p = 0.30$) over the decade; discharge from hospital alive rates have increased from 1.4% to 6.8% ($p = 0.017$) for octogenarians and not changed for nonagenarians (1.9%, $p = 0.82$). Survival to hospital discharge rates for octogenarians and nonagenarians in VF/VT were 10.3% and 2.5%, asystole 1% and 0.5% and PEA 3.4% and 3.4% respectively. Logistic regression analysis was used adjusting for Utstein data elements and using the 65-80 year old age group as reference. The odds ratio [OR (95% CI)] for survival to hospital discharge for octogenarians was [0.6, (0.47-0.77)] and for nonagenarians was OR = [0.45, (0.22-0.93)]. For octogenarians, having an initial shockable rhythm is the strongest predictor of survival to hospital discharge [OR = 5.12, (3.3-7.8)] and witnessed arrest [OR = 2.64 (1.58-4.41)] was also a positive predictor of survival.

Discussion

Survival to hospital discharge rates have improved in the last decade for octogenarians but not for nonagenarians. Quality of life measures are required in OHCA research to inform this debate.

OUT-OF-HOSPITAL CARDIAC ARREST IN YOUNG ADULTS IN MELBOURNE

**Dr Conor Deasy, MB, BAO, BCH, FCEM¹⁻³ Janet Bray, MCH² Dr Karen Smith, PhD
Dr Linton Harris, PhD A/Prof Steve Bernard, MD, FJFICM¹⁻³ Prof Peter Cameron, MD,
FACEM^{1,3}**

Introduction

Using the Victorian Ambulance Cardiac Arrest Registry (VACAR) we sought to determine the characteristics and outcomes of out-of-hospital cardiac arrest in the young adult population.

Methods

VACAR was searched for all out-of-hospital cardiac arrests (OHCA) occurring in young adult aged between 18 and 39 years, that were unwitnessed by EMS.

Results

Between 2000 and 2009 there were 30,227 cardiac arrests of which 3,820 were in this age group. The median age was 30 years for both sexes with a 3:1 male to female ratio. The most common precipitant was drug overdose (n=1299, 34%) followed by presumed cardiac (n=798, 20.8%) and trauma (n=718, 18.7%). Bystander CPR occurred in 18% overall, 27% where the precipitant was presumed cardiac and 36% where drowning was the precipitant. EMS median response time was 7 min, resuscitation was attempted in 35% of OHCA's. The presenting rhythm was asystole in 84%, PEA in 8% and VF/VT in 6%. Of those where EMS attempted resuscitation (n=1337), return of spontaneous circulation (ROSC) was achieved in 311 (23%) patients who were then transported to hospital, 106 patients surviving to leave hospital (8%). There were 138 patients (10%) transported with ongoing CPR of whom 7 (5%) survived to leave hospital. On multivariate analysis, more recent year of cardiac arrest was positively associated with increased likelihood of being discharged alive (OR 1.13, 95% CI 1.04-1.22, p=0.002). The survival to leave hospital rates for asystole, PEA and VF/VT were 1.5%, 10% and 31% respectively. Overall, the OHCA survival to leave hospital rate was 8% where EMS attempted resuscitation.

Discussion

Drug overdose is the most common cause of OHCA in young adults; preventive strategies are required to save lives.

LESSONS LEARNED FROM THE ‘ACADEMISATION’ OF THE HEALTH PROFESSIONS

Scott Devenish¹ MVEd, BNur, DipParaSc, RN, MACAP
Stephen Loftus² BDS, MSc, PhD

¹Lecturer Bachelor of Health Science (Paramedic), School of Public Health, Queensland University of Technology ²Deputy Director, Education for Practice Institute, Charles Sturt University

Background

Paramedicine is the healthcare provided by ambulance officers. At present, Paramedicine in Australia is gradually being ‘academised’, with the education of paramedics moving from industry into the universities (the academies). This is a trend established long ago, when other health professions moved into the universities, beginning with medicine and pharmacy. Over the years since then, other health professions have made the move to the university, such as dentistry, and more recently, physiotherapy and nursing. Now, all new entrants to these professions must begin by acquiring a university education.

Objectives

In this presentation we look at the lessons that can be learned from the ‘academisation’ of the health professions. These lessons can enlighten the current transformation occurring within the education of paramedics.

Methods

A review of the current literature was undertaken.

Findings

In the short term, all professions faced challenges such as industry demand outstripping university output, shortages in clinical placements, and pressure on existing staff to upgrade their qualifications along with a ‘generation gap’ between old and new. In the long term, ‘academisation’ offers the opportunity to improve professional status. Academic freedom can give a profession the chance to deliver a curriculum that reflects the reality of practice, and develop higher order critical thinking skills. Along with this comes the chance to develop a profession that is research-based. These days interdisciplinary health education can broaden the outlook of a profession within the wider society. These factors can promote lifelong learning and produce professionals who are willing and able to contribute to the knowledge base of their profession.

Conclusions

The challenges faced as a result of paramedic education moving to the universities are not new. As seen with other health professions, the long term benefits of ‘academisation’ can outweigh the difficulties in the short term.

THE PROFESSIONAL SOCIALISATION OF PARAMEDICS: UNIVERSITY AND ITS ROLE IN THE SOCIALISATION PROCESS

Scott Devenish¹ MVEd, BNur, DipParaSc, RN **Michele Clark**² BOccThy, BA, PhD
MaryLou Fleming³ DipTeach, BEd, MA OHIO S, PhD **Stephen Loftus**⁴ BDS, MSc, PhD

¹Lecturer, Paramedic Practice, School of Public Health, Queensland University of Technology, Brisbane Australia, ²Director of Research, School of Public Health, Queensland University of Technology, Brisbane Australia, ³Head of School, School of Public Health, Queensland University of Technology, Brisbane Australia, ⁴Deputy Director, Education for Practice Institute, Charles Sturt University, Sydney, Australia.

Background

Professional socialisation is defined as the process by which professionals learn the values, behaviours and attitudes necessary to assume their chosen professional role. The socialisation process consists of three distinct phases namely, pre-socialisation (Anticipatory), socialisation (Encounter) and post-socialisation (Change and Acquisition) phases.

Objectives

This research investigates the university's role in the professional socialisation of undergraduate paramedic university students by exploring the lived experience of students about to make the transition from university to practising paramedic. In this presentation, we explore the experiences of students moving from the anticipatory phase to the encounter phase.

Methods

Final year undergraduate university students from Charles Sturt University were approached to take part in this study. Data collection was via semi-structured interviews. A thematic analysis of the transcripts was employed using a hermeneutic phenomenological approach.

Findings

Prior to enrolling in a paramedic undergraduate university course, the majority of students interviewed believed paramedical work was mostly 'lights and sirens'. This view was predominantly formed by seeing or hearing ambulances rush to emergency cases, passing ambulances at car accidents, the television news and hearing stories told by paramedics. Television shows such as All Saints, however, seemed to have very little influence on their preconception. The motivating forces behind students wanting to pursue a career as a paramedic were varied. After beginning university studies in paramedical practice, information taught in classes and experiences on clinical placements was in conflict with their preconceived beliefs. Students had to undergo a detachment process from their former expectations.

Conclusions

The university system has an important role in supporting the professional socialisation of paramedics. Socialisation can be a very difficult experience and may be very confronting if there is discordance between preconceived expectations and reality.

SECONDLOOK: ENHANCING THE EDUCATION OF PARAMEDICS AND THEIR DYNAMIC RESPONSE TO PATIENTS WITH COMPLEX SOCIAL NEEDS

Kathryn Eadie^{1,3} (BBEHSc, HonsPsych) **Dennis Jess**¹ (Associate Diploma of Health Science (Amb), ASM) **Joanne Stephens**¹ (BNurs, DipHlthSc(Amb), MAdVocEd) **Monique Edmunds**¹ (BIntBus, BA AsIntSt), **Vivienne Tippet**^{1,2,3} (BA, Grad Dip Psych, MPH, PhD) **Leo McNamara**¹ (Associate Diploma of Health Science (Amb), ASM)

¹Queensland Ambulance Service, Brisbane, Queensland, Australia

²School of Population Health, University of Queensland, Brisbane, Queensland, Australia

³Australian Centre for Prehospital Research, Queensland, Australia

Objectives

The purpose of the Vulnerable Clients Program Initiative is to develop and deliver face to face paramedic education complemented by a self directed DVD package to increase awareness of the needs of patients with complex social needs. Passive referral tools have been developed for vulnerable clients and a communication board has also been developed for clients with communication difficulties. A rolling evaluation of the impact of the program on staff awareness is being conducted.

Methods

Primary data collection of quantitative and qualitative data will inform baseline, uptake and outcome analyses of the proposed education program. As social issues are not routinely coded in the electronic Ambulance Report Form (eARF), data on mental health which is more consistently collected will be used as a loose 'proxy' measure of QAS exposure to the vulnerable client target group.

Other descriptive information on the possible extent of ambulance service usage by vulnerable clients will be sourced from existing publically accessible government reports (e.g Australian Social Atlas, Australian Bureau of Statistics, Census, Office of Economic and Statistical Research). Data sources for evaluation of the educational package include the following:

1. Paramedics complete a pre-training survey comprising measures of awareness, attitudes & behaviour
2. Paramedics complete a post-training survey comprising training evaluation and confidence measures of utilizing tools
3. Paramedics complete a six month follow-up survey comprising measure of awareness, attitudes & behaviour, implementation of passive referral tools and a communication board
4. Paramedics report use of passive referral tool and communication board on eARF, and document vulnerable groups attended to on eARF

Findings and Conclusion

Baseline data on the incidence of vulnerable client groups in the pilot QAS Region will be reported, as well as a preliminary evaluation of paramedic training. Preliminary data will highlight the importance of the vulnerable client program training and the need for referral tools to assist patients with complex social needs.

PARAMEDIC ACCURACY IN IDENTIFYING THE ANATOMICAL LANDMARKS FOR NEEDLE CHEST DECOMPRESSION

Dale Edwards¹ Lisa Bowerman¹ Brett Gibson² Danielle Berry²

¹University of Tasmania, School of Medicine, Hobart, Australia. ²Ambulance Tasmania, Hobart, Australia

Background

Within the field of paramedicine, there are a number of clinical skills and interventions practiced on an infrequent basis, and have a significant impact on patient outcomes, these include such skills as chest decompression. This project will evaluate the capacity of paramedics at all levels of qualification and experience to accurately identify the anatomical landmark for insertion of a chest decompression needle.

Objectives

To provide evidence regarding the accuracy of paramedics in locating the anatomical landmarks required for the anterior placement of a needle for chest decompression. The sample was opportunistic, making use of attendees at the ACAP Tasmanian state conference in June 2010

Methods

Following the identification of the true mid clavicular line on a patient role player, utilising ultrasound to locate the true distal point of the clavicle, and subsequent 2ICS MCL, a removable mark is placed on the patient. This mark is captured by digital image as a reference point, then removed. Participants of the study were asked to identify on the role player the 2ICS MCL, and place a mark. Each participant's mark was recorded, then removed in preparation for the next participant. Participants were also asked to complete a brief survey to provide contextual data.

Digital images will be analysed for measurements utilising *ImageJ*. Data gathered will be analysed using a one way t test as well as through correlation of survey responses to accuracy findings.

Findings

This project is ongoing, with data analysis yet to be completed. This will be completed prior to the conference, and ready for presentation.

Conclusions: As Above

KEEPING PARAMEDICS SAFE: EXPERIENCES IN DELIVERING A “SELF DEFENCE” PROGRAM TO QUEENSLAND PARAMEDICS

Helen Eyles, Brian Pink

Queensland Ambulance Service, Queensland, Australia

Background

In recent time, a number of Queensland Ambulance Service (QAS) paramedics have been assaulted or their personal safety has been placed at risk in the performance of their normal duties. A project team was appointed in November 2009 dedicated to identifying aspects that contribute to prehospatal workplace violence in Queensland. The team was to use this information to design and deliver a course that could reduce the impact of workplace violence being experienced by front line paramedics.

Objectives

To establish a baseline level of behavior and tactics for managing workplace violence. Paramedics must be able to immediately apply course concepts regardless of previous experience or pre-existing fitness levels.

Methods

Course design was based on data generated from WH&S reports, the QAS Commissioner’s imperatives and a contracted content expert’s expertise.

Findings

Situational Awareness for Everyday Encounters (SAFE) is an 8 hour mandatory course for operational staff of the QAS. The rollout began in March 2009 and the teaching team has been tasked to deliver the course to all 3300 officers (across the State) by the conclusion of 2010.

With no Australian based, paramedic-centred course to use as a template, the team has had a unique opportunity to identify strengths and shortfalls in course content and delivery. The QAS is now developing a body of knowledge in the practical delivery of this type of course to share with interested paramedic services.

Conclusions

The SAFE program is another aspect of the Queensland Ambulance Service’s commitment to the “Zero Harm” policy, and falls within the Q2 Queensland Government strategic plan. Key performance indicators will, in time, provide objective data on the impact of the SAFE program. Anecdotally, participant feedback for this non-clinical program has been overwhelming positive. Paramedics have applauded the initiative to invest in service delivery safety.

FALLS IN THE PREHOSPITAL ENVIRONMENT – A REVIEW OF THE LITERATURE

Emma Flavell¹ BN, RN **Malcolm Boyle¹** ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClInEpi, PhD

Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

Falls occur throughout one's lifespan, but are much more frequent in children and in the elderly, with at times with devastating consequences. Falls can be caused by a simple as a loss of balance or a more serious sign of underlying pathophysiology. Falls in the prehospital environment range in severity and are associated with significant morbidity and mortality. Falls attended by Ambulance Victorian amount to approximately 45% of their workload in a one year period.

Objective

The objective of this study was to identify the incidence of falls and the outcome from an ambulance perspective.

Method

A literature search was conducted using electronic databases Ovid Medline and EMBASE. These databases were searched from January 1996 to the end of May 2010. All references from the articles retrieved were also reviewed. Articles were included if they reported on falls attended by an ambulance crew.

Results

A total of 406 articles were identified. Of these, 30 articles met the inclusion criteria. However, only two articles were based in Australia (one in combination with New Zealand), with the remaining 28 articles from the U.S.A., United Kingdom, Scotland, Turkey, Iran and Pakistan. The literature found was based on the presentation of falls in the emergency department after being brought in by ambulance. These articles reported the frequency of falls presenting at the emergency department, from 4% to 34%, and highlighted the significant morbidity and mortality, 34%, that can arise from falls, particularly in paediatrics and the elderly.

Conclusion

Despite the relative frequency of falls there is a substantial gap in the prehospital literature and the implications for paramedic practice. There is a need for more research into the incidence of falls attended by ambulance and the patient's outcome.

WHICH IS MORE EFFECTIVE FOR VENTILATION IN THE PREHOSPITAL SETTING DURING CARDIOPULMONARY RESUSCITATION, THE LARYNGEAL MASK AIRWAY OR THE BAG-VALVE-MASK? - A REVIEW OF THE LITERATURE

Emma Flavell BN, RN Malcolm Boyle ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClinEpi, PhD

Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

Prehospital care providers are responsible for providing adequate ventilation during cardiopulmonary resuscitation (CPR). Endotracheal intubation (ETI) is widely accepted as the ‘gold standard’ for airway protection and the preferred method for ventilation. However, most paramedics are not trained to perform ETI. Laryngeal Mask Airway (LMA) and Bag-Valve-Mask (BVM) are seen as adequate alternatives to ETI as recommended by the International Liaison Committee of Resuscitation (ILCOR).

Objectives

The objective of this study was to identify which airway device LMA or BVM (with OPA/NPA) is more effective in airway patency and ventilation during cardiopulmonary resuscitation in the prehospital environment.

Methods

A literature search was conducted using medical electronic databases, MEDLINE CINHAL, EMBASE, Meditext, Cochrane Central Register of Controlled Trials (CENTRAL), and Scopus. These databases were searched from January 1996 until the end of December 2009. References from articles retrieved were reviewed. Articles were included if they reported a comparison of LMA and BVM with humans or mannequins for the prehospital setting.

Findings

There were 2,937 articles located by the search. Of these, 30 articles met the inclusion criteria with twelve relevant to the prehospital environment. In the twelve prehospital studies there were two mannequin, four retrospective, and five observational studies, with one literature review. The studies concluded that although the LMA may take longer to initiate the first ventilation; it provides more effective ventilation over a longer timeframe and decreases gastric regurgitation and inflation, especially in adults. Evidence of LMA use in paediatrics is inconclusive for the prehospital setting.

Conclusions

Findings from this review suggest that the LMA is more effective at ventilations during CPR in adults, as there is less risk of gastric regurgitation and pulmonary aspiration. The BVM is quicker at performing the first ventilation but there is a loss of effectiveness over time. BVM is still considered the best method for ventilating children and neonates.

DOES EVERY SECOND COUNT? A LITERATURE REVIEW OF THE EFFECT OF AMBULANCE RESPONSE TIMES ON PATIENT OUTCOMES IN CASES OF TRAUMA

CoCo Giddings, Nicole Robertson, Malcolm Boyle, Brett Williams

Department of Community Emergency Health and Paramedic Practice, Monash University,
Melbourne

Background

Ambulance response times are prioritised by ambulance management and receive a significant amount of attention from government agencies and the broader community. Ambulance legislation frequently includes guidelines for response times and are utilised as key performance indicators on the presumption that patient care and outcome is improved if ambulance response time is shorter. The available data suggests that whilst this is certainly the case for patients in non-traumatic cardiac arrests, little attention has been paid to the effect of ambulance response times on patient outcomes in other health care emergencies.

Objectives

The objective of this study was to examine the existing evidence surrounding the effect of ambulance response time on patient outcome in cases of trauma.

Methods

A review of selected electronic databases was conducted from their commencement to the end of May 2010. Inclusion criteria were any study type reporting on the relationship between ambulance response times and trauma patient outcomes. References of relevant articles were also reviewed.

Findings

The search located approximately 180 articles and 14 articles met the inclusion criteria. The limited number of studies conducted has produced contrasting results, with one study showing strong evidence to support the argument that response times have little impact on trauma patient outcomes, whilst others with lower levels of evidence indicate that shorter response times produce better outcomes in trauma patients. Based on these levels of evidence and the power of the results achieved in each of the studies it would appear that stronger support is present for the argument that response times do not predict patient outcomes in trauma cases.

Conclusion

The available literature questions the link between response times and trauma patient outcomes. There is evidence to suggest that shorter response times do not produce better outcomes, and that alternate performance indicators may be more appropriate in these cases.

SOUTH AUSTRALIAN AMBULANCE SERVICE (SAAS) EXTENDED CARE PARAMEDIC (ECP) PILOT PROJECT

Hugh Grantham^{1,2} ASM, MBBS, FRACGP
Cindy Hein^{1,2} DipAppSc, BHSc, PhD **Rob Elliott**^{1,2} BHSc, GradDipBA

¹SA Ambulance Service (SAAS), Adelaide, Australia ²Flinders University, Adelaide, Australia

Background

The SAAS ECP pilot project (1st December 2008 – 30th June 2009) was instigated as a response to a review of the current and predicted ambulance workload within SAAS. This review indicated the need to explore strategies that change the way of delivering ambulance services from a traditional emergency response/emergency department model, to a broader approach inclusive of dynamic clinical care.

Objectives

This small scale preliminary study was established to pilot ECPs working with other health professionals with the aim of providing initial assessment and treatment of patients in their own homes or residential care facilities, as opposed to routinely transporting them to an emergency department.

Methods

Extensive liaison with southern Adelaide health services occurred over many months prior to launching the pilot, thus establishing a multidisciplinary quality assurance (QA) team. 12 intensive care paramedics were selected and trained for the role and specially designed/equipped vehicles were purchased. Four key performance areas measured included: 1) QA of each patient management; 2) Patient feedback; 3) General Practitioner (GP) feedback; and 4) activities of the ECP clinician role in the Emergency Operation Centre (EOC).

Findings

ECPs attended 1123 patients, of those 555 interventions (49.4%) were considered to have prevented an ED presentation and 60 (5.3%), were considered to have prevented a hospital admission. The mean, median (range) age was 71, 78 (14-102) years respectively. No adverse events were recorded. ECPs in the EOC provided 2815 self-reported qualitative reports from 7 activities (refer Table 1). Feedback from patients and carers, as well as doctors, was extremely positive.

Conclusions

This pilot addresses not only the needs of the ambulance service and the health system, but also the needs of the patient and was supported by health services as part of a suite of strategies to address the workload issues found in the public health system.

UNDERGRADUATE PARAMEDIC CPR FATIGUE: RESULTS FROM A PILOT STUDY

Hendrik Gutwirth¹ (BAppSc (Exercise and sport), BAppSc (sport science -Tennis), MAppSc (Strength and Conditioning), Exercise Physiologist (AAESS) **Brett Williams**¹
Malcolm Boyle¹ **Trevor Allen**²

¹Department of Community Emergency Health and Paramedic Practice, Monash University

²Department of Physiology, Monash University

Background

The 2005 ILCOR guidelines suggest rescuers deliver CPR in cycles of 30 chest compressions and 2 ventilations (30:2) at a rate of 100 compressions/minute with a compression depth of 4-5cm. Given this increase from the previous CPR cycle, 15:2, there is now greater emphasis on pushing faster and deeper with minimal interruption. This has led to speculation surrounding rescuer fatigue and compression efficiency.

Objectives

The objective of this study was to identify the level of fatigue and the quality of chest compressions during simulated CPR.

Methods

This was an observational pilot study investigating second year undergraduate paramedic students' fatigue levels and quality of chest compressions following twenty minutes of simulated CPR on a Laerdal Resusci Anne mannequin. Data were collected at baseline and every 2 mins until conclusion of twenty minutes. Student heart rates were collected using a wireless heart rate monitor with student fatigue measured using the Borg Scale.

Findings

Seven students participated, two were males with five students between 21 and 25 years, the others were <21 years. There was a statistically significant difference between males and females for compression rate, 108.0 Vs 125.6 chest compressions/minute, $p=0.006$. There was a statistically significant difference between base heart rate (resting prior to commencing CPR) and heart rate at 14 minutes, $p=0.045$, for all students and for the Borg Scale at rest and at 2, 6, 10, 14, and 18 minutes, ($p<0.01$). There was a statistically significant difference between the <21 year group and 21-25 year group for compression rate, mean 113.4 Vs 136.1 chest compressions/minute, $p<0.001$.

Conclusion

This pilot study suggests that fatigue sets in early when undertaking CPR in a controlled setting with some fatigue attributed to the faster than recommended chest compression rates. Further research is required to identify if student fitness levels affect the rate of onset of fatigue.

MENTAL HEALTH CASES ATTENDED BY AMBULANCE PARAMEDICS IN METROPOLITAN MELBOURNE USING ELECTRONIC CLINICAL DATA

Linton Harriss, Karen Smith, Alex Currell

Ambulance Victoria, Melbourne, Australia

Background

One in five Australians will experience mental illness at some stage in their life, and for many of these people, paramedics represent the first contact with the health care system. Previous analysis of mental health attendances by Ambulance Victoria (AV) was limited by identification of patients via Medical Priority Dispatch System (MPDS) codes. These codes are allocated at the call-taking stage where the condition of the patient may differ to that actually found in-field. The introduction of the Victorian Ambulance Clinical Information System (VACIS), allows for identification of mental health cases via electronic clinical data gathered by paramedics. This improves sampling and analysis.

Objectives

To conduct a comprehensive epidemiological evaluation of mental health cases attended by AV paramedics in metropolitan Melbourne using VACIS data.

Methods

A retrospective cohort was used to investigate characteristics and care of mental health cases attended from 1 July 2008 – 30 June 2009. Cases were classified according to involuntary status, transport/treatment status and presentation.

Findings

A total of 22,885 mental health cases (13,184 women) with mean age of 43.5 years (SD, 19.8) were attended by paramedics for the one-year study period. This represents approximately 8.0% of annual AV emergency attendances. The majority of cases required either no transport or transport only (80.4%) and had a documented mental health history (75.9%). Relatively few cases were involuntary (7.4%), had a mental health MPDS code (34.7%), involved CATT (10.0%) or Police (18.1%) at scene, or required treatment (21.1%), restraint (2.4%) or medication (6.2%). Most common presentations were psychosis (30.8%), social emotional problems (15.3%), substance related (10.4%), anxiety (9.0%), and mood disorders (8.7%).

Conclusions

The majority of mental health cases attended by paramedics don't require specialist care. Results from this research will assist in enhancing AV service delivery to these patients and improve access to appropriate health care facilities.

EXTENDED ROLES, EXTENDED SCOPES, ACPs, ECPs, WHAT'S IN A NAME? AN EXAMINATION OF THE NON-TRADITIONAL ROLES OF THE PARAMEDIC IN AUSTRALIA, UNITED KINGDOM, UNITES STATES AND CANADA

Peter Hartley MHIthSc Vic(Aust), CertIVWorkTrain&Assess,
CAECA, GradDipEd (Melb.), DipHIthSc(AmbOfficer),VPSEB, AssocDipSocStuds

School of Biomedical and Health Sciences, Victoria University, Melbourne, Australia

Background

As most outer lying communities throughout the western world struggle to attract local medical services, the role of the paramedic is increasingly being looked upon to assist in the provision of health care to the community it serves. This has necessarily required that EMS providers across the world have been required to pursue innovative models of service delivery to meet community defined needs. Collaboration of EMS and community organizations such as primary health care providers, social service agencies, and public safety groups have enabled innovative initiatives that have now proven to improve the level of health care within a community, both in rural and urban settings.

Many models have been implemented overseas, and have incorporated expansive programs ranging from supporting volunteers and treating patients in community as opposed to transporting. The models have addressed such issues as the expanded role of the community paramedic, paramedics without borders, and such complex issues pertaining to the balance between rural versus urban settings and rural and remote issues in community paramedicine.

There is no doubt that the extended scope of practice for paramedics in Australia has become a contemporary issue, as it has done worldwide. Only the varying models being disputed have compounded the complexities of this issue.

Objectives

The paper will report on the author's visit to varying models of community paramedic programs throughout the United Kingdom, Unites States of America and Canada, and offer a comparative analysis to Australian models and needs.

Methods

Data gathered from physical site visits to programs throughout comparative nations, with specific emphasis on the elements that form a robust community focused program.

Conclusions

This paper presents the comparative programs across four nations as they relate to the non-traditional role of the paramedic, and will offer conclusions as to why some are seemingly more robust.

PARAMEDIC HIGHER EDUCATION, WHO LEADS, WHO FOLLOWS AND WHO COLLABORATES? AN EXAMINATION OF THE ROLES, FUNCTIONS AND CHALLENGES FACING PARAMEDIC EDUCATION IN THE AUSTRALIAN HIGHER EDUCATION SECTOR

Peter Hartley MHLthSc Vic(Aust), CertIVWorkTrain&Assess, CAECA, GradDipEd (Melb.), DipHlthSc(AmbOfficer), VPSEB, AssocDipSocStuds Assoc ACAP AERA

School of Biomedical and Health Sciences, Victoria University, Melbourne Australia

Background

Paramedic education in Australia has encountered a considerable evolution this past decade. As EMS providers are faced with increasing public demand for services, the paramedic education system is equally challenged to respond to such needs, whilst in many ways is still 'cutting its teeth' in the realms of academia. This environment necessarily requires that consistently improved models of paramedic education are essential to ensure that graduate paramedics are prepared for the ever-increasing demand for more complex skills sets and growing requirements for autonomy in practice.

Authorities demand that paramedic higher education programs produce work-ready graduate paramedics with not only appropriate clinical ability but also a multi skilled practitioner. This in essence can be in contrast to the higher education (HE) mandate of university graduates. As the paramedic HE system struggles to establish itself in the dominion of academia, coupled with the complexities and contrasts of its established roots in vocational education, the major challenge emerging appears to be the definitions and expectations of graduates.

Objectives

Higher education places great emphasis on developing graduates as critical thinkers, to question beyond the expected, to challenge and forge beyond the conventional. It is no surprise then when conflict arises as graduate paramedics enter a working environment that is steeped in specific organizational guidelines and instructions. The question that fundamentally arises from this is one that investigates the appropriateness of critical pedagogy in higher education paramedic programs.

Further to these challenges are the multifaceted ideologies formed from basic underlying principles of who leads, who follows, and who collaborates.

Methods

Systematic review of literature.

Conclusions

Challenges continue to face higher education paramedic programs in Australia and conflicts continue to arise from the transition from VE/FE. Equally, conflicts face graduate students as they themselves make the transition from HE institutions to the paramedic workforce.

SAVING LIVES AND PREVENTING INJURIES BY REDUCING AMBULANCE VEHICLE CRASHES - THE ROAD SAFETY BLACK BOX SYSTEM FOR AMBULANCE FLEETS

Peter Hartley¹ MHLthSc Vic(Aust), CertIVWorkTrain&Assess,
CAECA, GradDipEd (Melb.), DipHlthSc(AmbOfficer),
VPSEB, AssocDipSocStuds Assoc ACAP AERA
Jerry Overton²

¹ School of Biomedical and Health Sciences, Victoria University, Melbourne, Australia ²
Road Safety International, Thousand Oaks California, United States of America

Background

Much has been reported over the years as to the hazards involved for both paramedics and patients alike in respect of ambulance transport. There is significant evidence that supports statements that in emergency responses, increases in braking and acceleration have considerable impacts on patient treatment regimes as basic as the effectiveness of CPR. Sixteen years ago, Road Safety International brought the "Black Box" from the airplane cockpit to the highway. The mission was to insure the safe operation of fleet vehicles by eliminating unsafe driver behaviors using on-board monitoring and instant feedback systems. When fleet management focuses on safety, other critical areas such as vehicle availability, maintenance and operational costs realize tremendous benefits. The Road Safety System is the most advanced, cost effective and easy to use tool for improving safety and reducing operating costs available to ambulance fleets.

Road Safety is the market leader in driver safety and performance monitoring systems. The SafeForce™ Driving System provides driver monitoring, real-time feedback, data analysis and driver performance reporting. The System's real-time driver feedback and reporting information have produced dramatic crash reductions and cost saving results.

Objectives

Drivers of ambulance vehicles sometimes confuse getting on-scene quickly and safely with getting there at any cost. The result is intersection crashes, rear-end collisions and backing incidents, many of these incidents are predictable and preventable.

Methods

This paper uses a systematic review of white papers and documents relating to the effect of the SafeForce™ Driving System and will utilize real-time electronic data from on board computer records in ambulance vehicles over a 24-month period.

Conclusions

The SafeForce™ Driving System is a proven method of changing driver behaviors and dramatic improvement in road safety of emergency ambulance vehicles.

**IT'S THE END RESULT THAT COUNTS...PERHAPS!
A COMPARATIVE REPRESENTATION OF PARAMEDIC GRADUATES FROM
VICTORIA UNIVERSITY, MELBOURNE, AUSTRALIA DEGREE PROGRAM, AND
THE COLLABORATIVE EDUCATION APPROACH IN MD AMBULANCE
SASKATOON CANADA**

Tim Hillier¹ Cert A&CE, Sask, ACLS, BCLS, BTLS, PEPP, CAMATA (Aeromedicine)

Peter Hartley² MHLthSc Vic(Aust), CertIVWorkTrain&Assess,
CAECA, GradDipEd (Melb.), DipHlthSc(AmbOfficer),
VPSEB, AssocDipSocStuds Assoc ACAP AERA

¹MD Ambulance, Saskatoon Saskatchewan Canada, ² School of Biomedical and Health
Sciences, Victoria University, Melbourne Australia

Background

Paramedic education in Australia has continued to evolve and embrace the concepts of higher education for the last decade. This has brought about a considerable move from its established roots in vocational teaching and seen a significant shift from the previously collaborative and service owned approach to paramedic edification to the now more autonomous programs that exist in higher education degrees, often with comparative limited clinical exposure. Questions continue to be asked as to the impact this has on graduate attributes and the work readiness of the potential workforce for EMS providers. This is a comparative presentation of the attributes of Victoria University graduate students' knowledge and skill sets with those of Advanced Care Paramedics employed by MD Ambulance, Saskatoon.

Objectives

To determine if there is any dissimilarity in the work readiness of paramedic students between one university higher education degree program in Australia and that of a paramedic education program that has a strong collaborative emphasis with an EMS provider in Saskatoon Canada.

Methods

Narrative and systematic review of core graduate attributes, curriculum and skill sets and comparative analysis of group 1 Victoria University graduate students, and group 2 Advanced Care Paramedics employed by MD Ambulance, Saskatoon Canada.

Conclusions

The delegation of paramedic education from the vocational sector to the more autonomous higher education programs continues to gather popularity as the pathway of the future. Whilst there appears to be some disparity between expectations of EMS providers and those of the university in relation to graduate attributes, such disparity is less so for programs that have a collaborative inclusive approach to education of paramedic requirements with the EMS providers.

DO EXTENDED CARE PARAMEDICS REDUCE HOSPITAL EMERGENCY ATTENDANCES?

Sarah Hoyle¹ MHLthSc, **Andy Swain^{1,2}** BSc, MBBS, PhD, FRCS, FCEM, FACEM
Peter Larsen² BSc(Hons), PhD

¹Wellington Free Ambulance, New Zealand

²Department of Surgery & Anaesthesia, University of Otago, Wellington

Background

In May 2009, Wellington Free Ambulance introduced extended care paramedics (ECPs) who had been trained in additional clinical skills to treat patients in their own homes or communities. The aims of this new model of care were avoidance of unnecessary transfer to hospital, improved patient experience, and more effective allocation of emergency ambulances.

Methods

In this prospective observational study we examined the first 12 months of the ECP programme to determine what types of patients had been seen by this service, and what proportion of these had been treated at home, rather than transported to the emergency department compared to standard ambulance crews.

Findings

Over a 12 month period, ECPs attended a total of 1351 cases. 61% of patients were female, with 7% under 16 years, 32% aged 17 to 64, 13% aged 65 to 75 and 48% over 75 years old. In total, 61% were treated at home or in the local community. This compares with a treat-at-home rate of 26% by standard paramedics for the same region and within the same time period. 7% of ECP cases had cardiac problems, 6.3% respiratory, and 22% were faints, near-faints or falls. Treatment at home or in the community did not differ significantly with patient age, but it did differ on the basis of the presenting complaint, with 44% of cardiac presentations, 55% of respiratory problems, and 40% of faints/near faints/falls treated at home ($p=0.001$, Chi squared test). Analysis of hospital records to determine whether any of the patients treated at home presented to emergency departments within 72 hours of ECP attendance is currently being undertaken.

Conclusion

This study suggests that ECP's have significant potential to reduce hospital emergency attendances, although this will differ depending on the presenting condition. This study provides justification for a randomised controlled trial comparing ECP with standard care.

CLINICAL DECISION MAKING DURING MASS-CASUALTY TRIAGE: A PRE-HOSPITAL PERSPECTIVE

Matthew Humar BAppSc (Human Movement) **Erin Smith** PhD

Department of Community Emergency Health and Paramedic Practice, Monash University,
Melbourne, Australia

Background

Paramedics must be able to make rapid and accurate triage decisions regarding the patient's criticality, the commencement or withholding of immediate care, and the need for extra resources at the scene of a mass casualty incident. Due to the infrequency of mass casualty incidents, the opportunity to practice and develop these triage decision-making skills is rarely afforded. Effective education and training are imperative within this area to better prepare paramedics' when making triage decisions at mass casualty incidents.

Objectives

The objective of this paper is to analyse the literature relating to decision-making during mass casualty incidents and to make recommendations to improve education and training methods within this area.

Methods

A review of the Ovid MEDLINE database (1950 to April 2010) was performed to locate relevant literature. Key search terms included: triage; protocol; algorithm; decision-making; pre-hospital; paramedic; mass casualty; multi casualty and disaster. Additionally, reference lists from relevant papers were reviewed to identify further relevant literature.

Findings

Triage is regarded as the most important function to be performed by paramedics at the scene of mass casualty incidents. Mass casualty triage decisions by paramedics are influenced by triage protocols, experience, stress, uncertainty, education and training.

Conclusions

Experience and/or building expertise is the principal factor influencing a paramedic's ability to make decisions at mass casualty incidents. The research directly concerning the role of paramedics in this field is limited and is an area that therefore requires future attention. Further research into this field is required to increase understanding and provide information that will serve as a basis for education and training programs.

PARAMEDIC INITIATED PAEDIATRIC RAPID SEQUENCE INTUBATION SUCCESS AND ITS ENDURING COMPLICATIONS

Jason Hunter

MICA Flight Paramedic, Air Ambulance Victoria

Rapid Sequence Intubation (RSI) by paramedics has long been a controversial issue. Between 2004 and 2008 Dr Stephen Bernard conducted a randomised controlled study in Melbourne on the effectiveness of pre-hospital rapid sequence intubation on traumatically brain-injured adults. Prior to this study MICA Flight Paramedics at Air Ambulance Victoria had been practicing paediatric RSI since 2000.

Between January 2006 and July 1 2010 MICA Flight Paramedics rostered to Air Ambulance Victoria's HEMS 1 and 5 base conducted 51 Paediatric RSIs. This number excludes any individual intubated without the use of pharmacological measures or was conducted with the assistance of a medical practitioner. All patients were transported to the Royal Children's Hospital, Melbourne or Monash Medical Centre, Clayton. The average age for those intubated was 7.7 years; they ranged from 5 weeks to 15 years. The average pre – intubation Glasgow coma score was 7.5. Importantly there were no recorded failed intubations within the case set. The predominant theme behind the paramedic initiated RSI was traumatic brain injury. The remainder included burns, respiratory failure, status epilepsy, non-traumatic head injury and post drowning.

Paediatric RSI is not without its complications, with haemo-dynamic compromise and hypoxia the most commonly documented. However one that is often not considered are the emotional effects tied to the case where paediatric RSI is required. Despite the average length of experience of a MICA Flight Paramedic being in excess of 10 years. Paediatric major trauma cases are often the most emotionally taxing cases one will encounter.

SCARRED FOR LIFE - AMBULANCE VICTORIA'S AERO-MEDICAL RESPONSE TO BLACK SATURDAY

Jason Hunter

MICA Flight Paramedic, Air Ambulance Victoria

The summer of 2008/2009 has been quoted by some as the most extreme ever encountered by Victorians. The forecast for February 7th 2009 was well publicised, temperatures were expected to be well above 45°C, low humidity and a hot northerly wind with a late change meant gusts in excess of 90 km/hr. It was to be business as usual for Ambulance Victoria (AV). With weeks of oppressive temperatures, the heat wave had placed extreme strains on the pre-hospital service. The AV emergency management plan was in action, the extra road resources deployed were already at breaking point.

Trouble was simmering days before with numerous fires being perilously close to breaking containment. The temperature in Essendon airport had peaked at 46.2°C. Air 9495 (HEMS 1) a SA365 N3 Dauphin helicopter was grounded, its capability in such extremis was beyond its current performance. The cool change swept across Melbourne late in the afternoon, 495 was back on line. The first request for assistance was received shortly after 1700 hrs. The Police Air Wing and 495 were tasked to winch up to 7 persons trapped by encroaching fires between Whittlesea and Kinglake.

This was only the beginning. By 1800 hrs all of Victoria's HEMS were airborne and were responding to tasks related to the bushfires. Throughout the 14-hour night shift the three MICA flight Paramedics had flown countless sorties from SAR to patient transport. More importantly, they had treated and assessed countless others and worked as integral part in one of Australia's worst natural disasters.

This was the allusive "big one" that all paramedics talk about. None of those involved could comprehend the mass of death and destruction that had occurred by shifts end. Patients and paramedics, scarred for life by the events of "Black Saturday".

1. David Parkham, 2009. Submission to Victorian Bushfires Royal Commission. May 2009. Sited at: <http://www.royalcommission.vic.gov.au/>.

NAMED PARAMEDIC MENTORS INCREASING GRADUATE ATTRIBUTES: THE PLYMOUTH UNIVERSITY EXPERIENCE

Annie Jenkin¹ RN, MSc, PGDE, BSc (Hons), HEA Fellow **Pam Nelmes**² RN, MSc, PGCE, BSc (Hons), Dip Nursing

¹School of Health Professions, University of Plymouth, Plymouth, United Kingdom. ²School of Nursing and Midwifery, University of Plymouth, Plymouth, United Kingdom.

Background

Higher Education based paramedic programs continue to face significant challenges; the most considerable of all continues to be those related to constraints of clinical exposure. With limited clinical experiences available, universities internationally are endeavoring to seek innovative approaches to ultimately increase the benefits of clinical experiences, and the graduate attributes for students in their programs. Paramedic courses at Plymouth University offer a unique Named Paramedic Mentorship (NPM) Program and are the only paramedic programmes nationally to allocate a specific mentor for the full 3 years.

Objectives

The objective of this program was to establish a clinical learning environment to provide paramedic students' maximum erudition within the constraints of limited exposure, and increasing graduate attributes.

Methods: The Named Paramedic Mentor approach is in its introductory and evaluative phase. To date, the first two student cohorts and equivalent number of Named Paramedic Mentors (N=72) are being evaluated and preliminary findings indicate significant benefits and improvements in the overall student experience and graduate attributes.

Inclusive in the evaluation are the supportive ancillary programs inclusive of mentorship feedback meetings, tripartite conferences, and mentorship training sessions.

Conclusions

This paper will present the Plymouth experiences to date, incorporating evidence to support this pedagogical approach. It will share best practice, the challenges and the strengths of embarking on having NPM for the duration of a student paramedic's Higher Education experience.

EVALUATION OF INDIVIDUAL AND ORGANISATION CLINICAL PERFORMANCE USING THE CLINICAL AUDIT AND REVIEW TOOL (CART)

Evan Lloyd MBA, BHsc¹ Sean Mutchmor BA, BSc, MA, CCP, ICP²

¹Queensland Ambulance Service, Principal Information Support Officer

²Queensland Ambulance Service, Manager, Clinical Standards

Background

One of the foundational aspects of clinical excellence is the capacity to accurately and consistently evaluate clinical practice.

Evaluations of clinical data have long suffered incompatible and unreliable evaluation with guidelines frequently inconsistent between jurisdictions. Consequently, it has been impossible to offer reliable data on clinical performance. Clinical improvement has relied on assessments based upon ad-hoc or isolated events; focused on causative factors rather than holistic performance.

Objectives

The development of the CART provides an application that interfaces with clinicians data capture. Outputs benchmark individual and organisational standards, articulate excellence and areas of concern providing evocative data used in developing clinical standards.

Methods

Electronic data was captured and distributed to clinical leaders via CART. Performance scores were calculated against established clinical measures. Complementary objective assessments contributed to an overall rating of clinical performance and returned to the attending clinician. Specific clinical triggers were assigned to organisational and patient care high risk activities.

Findings

The QAS now accurately identifies variables in clinical assessment / treatment, influencing both individual development programs and statewide education. There has been 1.08% improvement in quality care at variation levels 0 and 1 in the past 27 months. Levels 0 and 1 comprise over 98% of total evaluations. Clinical Support Officers have empiric evidence to design training based on need not assumptions. High risk cases are identified, evaluated and actioned proactively.

Conclusions

QAS relies on empiric data to determine clinical performance. Historical paper-based systems do not permit reliable reproducible systems of data analysis and performance measurement. CART has proven itself as a reliable, equitable platform that provides an indiscriminant, industrially friendly assessment tool. Clinicians now benefit from performance evaluation through an established set of criteria, benchmarks are well established and realistic, and clinical expectations are met and often exceeded through a constructive evaluation and feedback system.

ACCELERATED DEMAND FOR EMERGENCY AMBULANCES BY THE ELDERLY IN MELBOURNE: 1995-2008

Judy Lowthian¹ MPH, BAppSci (SpPath), LMusA **Damien Jolley**¹ MSc (Epidemiology), MSc, DipEd, A Stat, **Peter Cameron**¹ MBBS, MD, FACEM **Andrea Curtis**¹ BSc (Hons) PhD **Alex Currell**^{1,2} BEng, BA, MEngSc **Just Stoelwinder**¹ MD, FRACMA, FACHSE **John McNeil**¹ MBBS, PhD, FRACP, FAFPHM

¹Department of Epidemiology & Preventive Medicine, Faculty of Medicine, Nursing & Health Sciences, Monash University, Melbourne, Australia

²Ambulance Victoria, Melbourne, Australia

Background

Escalating demand places strain on emergency resources, with ageing suggested as a stress.

Objectives

We aimed to measure the growth in emergency ambulance utilisation since 1995 across Melbourne and quantify the contribution of population ageing.

Methods

A retrospective study using data provided by Ambulance Victoria and population data. Trends were analysed using various regression models.

Findings

The rate of emergency transportations has risen across Melbourne since 1995, from 31.8 to 57.6 per 1000 persons in 2008. The adjusted average annual growth was 5.5%, an overall increase of 75% in transportations (IRR=1.75, 95% CI: 1.6-1.9, $p<0.000$).

Compared to 1995, the number of observed transportations in 2008 was 68% greater than expected, after adjusting for age and gender.

Analysis of age/usage ratios revealed patients aged over 85 years utilised a higher proportional share of all transportations. By 2008 this age group rose from 1.1% to 1.6% of Melbourne's population, yet accounted for a rise in usage from 8.4% to 13.6% of emergency transportations. Adopting 45-69 years as a reference, transportation of patients aged 70-84 years had a 4-fold rise, with an 8-fold increase in those aged over 85 years, after adjusting for age and gender over the period. In patients aged 70-100+ years, the rate increased by 3% for each year of increasing age.

Conclusions

We confirmed accelerating demand by patients aged over 85 years. This age group will make up an increasing proportion of the future population, which will impact on ambulance services if transportations continue at the current rate. We also identified an additional 68% source of increase not explained by population change; therefore accurate identification and modelling of additional underlying drivers is needed.

UNDERSTANDING THE CHALLENGES: THE FACTORS THAT DRIVE THE DECISION TO ENTER AND EXIT INTENSIVE CARE PARAMEDIC PRACTICE IN AUSTRALIAN AMBULANCE SERVICES

Paul D McFarlane, RN, DipHlthSc (Nursing), BTh (Hons), DipMin, MHSts (Loss & Grief)
Volunteer Chaplain, Ambulance Service of New South Wales, Australia

Background

Intensive Care Paramedics provide a high level of emergency care to the Australian public, yet little is known about the reasons they choose this career or the issues that cause them to move away from frontline clinical work into different roles within the ambulance service.

Objectives

To determine the factors that drive the decision to enter or exit Intensive Care Paramedic practice in Australian ambulance services.

Methods

Fifteen Intensive Care Paramedics drawn from three different jurisdictions were interviewed using semi-structured qualitative interviews. The participants represented paramedics at three distinct career stages: newly trained practitioners; currently serving Intensive Care Paramedics; and those who no longer serve in frontline Intensive Care roles. The interviews were thematically analysed and yielded common themes that illuminated the factors influencing career choices.

Findings

Data analysis revealed that most participants desired to provide the highest level of clinical care for their patients and reported high levels of job satisfaction. This satisfaction was eroded by fatigue caused by shift work, deployment factors, management issues, exposure to trauma and the burden of responsibility. Participants reported on the likelihood of continuing in a frontline Intensive Care role long term, and the factors effecting that decision. Three main motivations for entering Intensive Care practice were identified. These motivational groupings were described as: Career Carers, Challenge Chasers, and Career Changers.

Conclusion

Intensive Care Paramedics are highly motivated clinicians who enjoy contributing to better health outcomes for the community. In order to retain their services, employers should seek to address issues related to fatigue, deployment concerns and barriers to communication between managers and on-road staff. Staff should be regularly educated about psychological self-care and injury prevention. Employers should also seek to provide working environments that are stimulating, challenging and well supported.

INTERPROFESSIONAL PEER TEACHING: STUDENT MIDWIVES TEACHING CARE OF MOTHER AND BABY AFTER BIRTH TO STUDENT PARAMEDICS

Gayle McLelland, Jill French

Monash University, School of Nursing and Midwifery Melbourne, Australia

Background

Although unplanned births before arrival (BBA) to hospital remains low, the latest Australian Institute of Health and Welfare statistics show that the number has doubled in 2006 compared to 1991. It would be expected that the primary health profession to care for this mother would be the paramedics. Whilst, the Australian paramedic discipline has seen a remarkable change in education over the last decade, there remains very little change in the content of the normal care of the mother and baby after birth. Inter-professional and peer teaching is increasingly popular in health education. Using inter-professional and peer teaching principles, this study investigates the possibility of student midwives teaching student paramedics the care of mother and baby after birth. Workshops in the care of mother and baby after birth were facilitated by the 3rd year undergraduate midwifery students for 2nd year undergraduate paramedic students.

Objectives

The object of this study was to investigate the effectiveness of 3rd year midwives' inter-professional peer teaching to 2nd year undergraduate paramedic students.

Methods

The findings were generated from an initial evaluation completed by the student paramedics on the day of the workshop. These were then followed up by surveys distributed to both the paramedic students and the midwifery students.

Findings

The findings for this study are currently being analyzed and will be available at presentation. However preliminary evaluations from both student midwives and student paramedics valued the inter-professional learning experience significantly.

Conclusions

Inter-professional education and learning opportunities are a vital to the development of the health professional student. Peer teaching has been a successful strategy in medical education. This study combined the principles of both to investigate inter-professional, peer teaching between 3rd year undergraduate midwives and 2nd year undergraduate paramedics. Overall the workshops were very successful with both cohorts gaining valuable insight and appreciation of the context of the different professional environment.

PROFESSIONAL CONFIDENCE: A POWERFUL ENABLING MECHANISM IN THE DEVELOPMENT OF CLINICAL PRACTICE

Paula McMullen, PhD, MHPed, BSN, RN

Senior Lecturer, Bachelor of Paramedic Practice, School of Medicine,
University of Tasmania

Background: Whilst the advent of specialisation in nursing and simultaneous development of the Specialist RN has been recognised throughout nursing literature, there is a paucity of published research regarding the RNs' experiences and processes in achieving this level of specialist nursing practice.

Objectives: The objective of this study was to explore the learning and professional development of RNs working in specialty areas as they progress from novice to expert.

Methods: The study utilised a mixed method design drawing on both quantitative and qualitative research methods. The quantitative component investigated the influence of learner attributes (their approach to learning, strategic control of study and self-efficacy) on the participants' academic learning outcome (n=39). While the quantitative methods allowed measurement and analysis of causal relationships between the presage factors and academic outcomes, the qualitative method focused on processes taking place during the learning process. This paper reports on the qualitative component, a largely naturalistic inquiry utilising a case study approach. The case drew upon multiple sources such as academic assessment tasks, clinical performance appraisals, reflective journals, open-ended questionnaires, semi-structured interviews and field notes. The case study (n=7) allowed for the elaboration of the learning process, the RNs' perceptions of these changes, and how these changes reflect the development of the characteristics of a Specialist RN.

Findings: The findings from this research revealed three integral components involved in this transitional process. For RNs to function competently in critical care areas, they must acquire and utilize a specialist knowledge base, develop domain specific clinical nursing competencies, and develop '*Professional Confidence*' as a Specialist RN. Professional Confidence emerged from the qualitative data as a composite phenomenon at the core of the process in the development of the Specialist RN. *Professional Confidence* is a powerful enabling mechanism in the transitional process. Professional Confidence comprised of four distinct dimensions.

Conclusions: Although this study is based on the learning and professional development of RNs, the transferability of these findings to paramedic practice is conceivable. There is paucity of research on the learning and professional development of paramedics. Would the same results occur if the study was replicated in paramedic students undertaking tertiary or 'industry/vocational' study, or in the ongoing professional development of paramedics throughout Australia?

ACHY BREAKY MAKEY WAKEY HEART?

Brae McWhinnie BClinicalPrac (Paramedic)¹ **Jason Poposki** BAppSc, BAppSc (Hons)¹
Malcolm Woollard MPH, MBA, MA(Ed), Dip IMC, PGCE² **Lettie Rawlins** BSc, MB, ChB³
Graham Munro MHSM, BHSc¹ **Peter O'Meara** BHA, MPP, PhD²

¹School of Biomedical Sciences, Charles Sturt University, Bathurst ²School of Biomedical Sciences, Charles Sturt University, Bathurst, Australia / Pre-hospital, Emergency & Cardiovascular Care Applied Research Group, Coventry University, UK ³Birmingham University School of Medicine, Birmingham, UK

Background

Music has been recommended as an aid to improving chest compression quality, but research indicates the popular European tune '*Nellie the Elephant*' reduces the proportion of laypersons compressing at the correct depth.

Objectives

Compared to no music (NM), does listening to '*Achy Breaky Heart*' (ABH) or '*Disco Science*' (DS) increase the proportion of CPR-trained health professionals delivering compressions at the 100bpm recommended rate and 4-5cm depth?

Methods

Randomised cross-over trial recruiting at the 2009 ACAP conference.

Findings

Of 74 participants 50% were male; median age was 37; 35% were ICPs, 26% paramedics, 20% students, and 19% other health professionals. 54% had taken CPR training within one year. Mode and inter-quartile range (IQR) for compression rate were NM 105 (99-116); ABH 120 (107-120); DS 104 (103-107). Differences between-interventions were significant for NM vs. ABH and DS vs. ABH ($p < 0.001$) but not NM vs. DS ($p = 0.478$). Compression rates of 95-105 were achieved with NM, ABH, and DS for 26/74 (35%), 8/74 (11%) and 34/74 (46%) of participants respectively. Differences were significant for NM vs. ABH ($p = 0.0005$) and DS vs. ABH ($p < 0.0001$), but not NM vs. DS ($p = 0.256$). Relative risk for a compression rate of 95-105 for ABH vs. NM = 0.31, for DS vs. NM = 1.31 (not significant) and for DS vs. ABH = 4.25. The number needed to *harm* was 5 for listening to ABH vs. NM and 3 for ABH vs. DS. A high proportion of compressions were too deep (NM 86%; ABH 79%; DS 77%, differences not significant).

Conclusions

Listening to *Disco Science* whilst performing CPR did not increase the proportion of pre-hospital professionals delivering compressions correctly. Unsurprisingly, listening to *Achy Breaky Heart* had a negative effect. Disconcertingly, regardless of the nature or absence of musical accompaniment, the majority of participants did not compress at the recommended rate or depth.

INTENSIVE CARE FLIGHT PARAMEDIC ARTERIAL LINE INSERTION: HAVE SUCCESSFUL INSERTIONS INCREASED?

Ben Meadley^{1,2} B.App.Sci(HM), Dip.Paramed.Sci, Grad.Dip Intensive Care Paramed, Grad.Dip Emergency Health (MICA), Grad.Cert Emergency Health (Aeromed).

¹Air Ambulance Victoria, Melbourne, Australia, and ²Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia.

Background

MICA Flight Paramedics (MFPs) from Air Ambulance Victoria (AAV) are authorised to insert arterial lines (AL). A 2008 study showed that MFPs were successfully inserting AL in 55% of all attempts.

Objectives

The primary objective of this study was to establish whether MFPs have increased their success rates of insertion of AL in the period since the 2008 study. A secondary objective was to determine whether opportunities to insert AL have increased.

Methods

Data was gathered from patient care records (PCRs) from three of the five AAV helicopters and also the fixed wing aircraft from January 2009 to May 2010. Inclusion criteria were: a MFP-staffed flight, at least one attempt at AL insertion, successful or unsuccessful insertion, primary or secondary response. Exclusion criteria were: physician-staffed cases, if the PCR was unclear as to whether AL was attempted, or illegibility of PCR.

Findings

209 cases were deemed suitable for analysis. Successful AL insertion was achieved by MFPs in 80% (n=40) of eligible cases. This represents a 25% increase in successful AL insertions in 18 months. In the majority of cases (59.8%, n=125), the hospital had already placed an AL prior to transfer. This result is consistent with the 2008 study (56%, n=114). Although this percentage had not significantly changed, the number of cases where MFPs had opportunity to insert an AL had increased (n= 22 in 2007/2008 vs. n=88 in 2009/10).

Conclusions

Successful AL insertion by MFPs has increased significantly in the 18 months since the initial study. This could be attributed to increased opportunities to perform the skill, as well as having been exposed to the skill for a greater period of time. These results show that MFPs are able to perform critical care interventions.

PRE-SHOCK CPR: SHOULD WE DO IT?

Jennifer Melvin¹ Brett Williams² Malcolm Boyle²

¹School of Medicine, The University of Notre Dame, Fremantle, Australia, ²Department of Community Emergency Health and Paramedic Practice, Monash University, Frankston, Australia

Background

Many studies over the past decade have investigated delaying initial defibrillation to perform cardiopulmonary resuscitation (CPR) as it has been associated with increased rates of restoration of spontaneous circulation and/or survival. The Australian Resuscitation Council (ARC), Electrical Therapy for Adult Advanced Life Support Guidelines (2006) recognised that results of clinical studies assessing the effectiveness of cardiopulmonary resuscitation (CPR) before defibrillation rather than a strategy of immediate defibrillation were inconsistent. Since 2006, a number of studies have researched these procedures.

Objectives

The objective of this study was to determine whether CPR before defibrillation is warranted.

Methods

A literature review was undertaken using the electronic medical databases Ovid Medline, EMBASE, CINAHL Plus, Cochrane Systematic Review, and Medtext from their commencement to the end of May 2010. Keywords used in the search included; CPR, defibrillation, VF, EMS, EMT, paramedic, emergency medical service, emergency medical technician, prehospital, and ambulance. The keywords were used individually and in combination. Inclusion criteria were, any study type reporting the use of CPR on patients in VF before defibrillation in the out-of-hospital setting. References of relevant articles were also reviewed.

Findings

Of the 3,336 articles located, eight met the inclusion criteria. The results of these studies indicated conflicting results. All retrospective studies (n=5) indicated a benefit in performing pre-shock CPR on patients with VF for durations between 90 and 180 seconds. Conversely, all randomised control trials demonstrated no benefit from providing CPR prior to defibrillation compared to immediate defibrillation for Return of Spontaneous Circulation (ROSC), neurologic outcome and/or survival to hospital discharge. However, none of the studies reported evidence that CPR before defibrillation is harmful.

Conclusion

Conflicting evidence remains regarding the benefit of CPR before defibrillation; however, no study concluded it was harmful. Therefore, further investigations are required to determine whether CPR prior to defibrillation improves patient outcomes.

THE PREHOSPITAL EPIDEMIOLOGY AND MANAGEMENT OF SPINAL CORD INJURIES IN NEW SOUTH WALES: 2004 – 2008

PM Middleton¹, S Anand^{1,2}, SR Davies¹, J Middleton²

¹ Ambulance Research Institute, Ambulance Service of New South Wales

² NSW Agency for Clinical Innovation Statewide Spinal Cord Injury Service

Background

The prehospital management of patients with a potential spinal cord injury is of critical importance, as up to 25% of spinal cord damage occurs after the initial injury. Previous studies have identified both immediate and effective spinal immobilisation and minimisation of the time to definitive care in a specialised spinal unit as of primary importance.

In NSW, an average of one hundred people are admitted annually to one of the state's two acute spinal care units following a traumatically acquired spinal cord injury. The majority of these cases are treated and transported by ground and air resources of the Ambulance Service of New South Wales (ASNSW).

Objectives

This study aimed to describe the epidemiology and prehospital management of all patients admitted to NSW spinal care units following traumatically acquired spinal cord injury.

Methods

Data describing the epidemiology and spinal cord injury patterns of patients admitted to NSW spinal care units are collated in the NSW Spinal Cord Injury Dataset (SCID). This study extracted patient identifiers from the SCID and linked them to ASNSW patient datasets for the time interval January 2004 to June 2008.

Findings

Over 85% of patients eventually diagnosed with spinal cord injuries were treated at the scene of their injury according to ASNSW spinal injury protocols, including appropriate spinal immobilisation. For these same patients, the median time to definitive care in a spinal care unit was twelve hours, with over 60% undergoing multiple hospital transfers before reaching a spinal unit. In 75% of cases most frequent chief presenting complaint was recorded as spinal injury or back pain, followed by multiple trauma then isolated head injury. This collaborative research has the potential to inform strategic health priorities.

EPIDEMIOLOGY OF ATRAUMATIC CHEST PAIN PRESENTING VIA TRIPLE ZERO IN NEW SOUTH WALES

Paul Middleton¹ Sowmya Anand^{1,2} Jillian Patterson¹ Bridie Carr²

¹Ambulance Research Institute, Ambulance Service of New South Wales, ²NSW Agency for Clinical Innovation Cardiology Network

Background

Prompt responses to symptoms of chest pain are important, as it may represent a myocardial infarction or other acute coronary syndrome, for which early treatment might be life-saving. An understanding of the epidemiology of triple zero calls for such symptoms is therefore needed to provide information such as the description of who these patients are and where they come from, and to evaluate the provision of services for chest pain patients.

Objectives

The aim of this project was to describe the epidemiology of triple zero calls where the paramedic on scene operated with a working diagnosis of chest pain.

Methods

Records from our Patient Healthcare Record (PHCR) database were extracted for cases where the patient was treated by any chest pain protocol. Adult emergency cases were considered for the period 1 July 2008-30 June 2009, and linked with corresponding Computer Aided Dispatch data. Geographic information obtained from the responding vehicles was used to map the regional distribution of calls.

Findings

In the 12 months to 30 June 2009, ASNSW responded to 46,722 calls where the paramedic treated the patient with a chest pain protocol. The median age was 69, 53% of calls were regarding males, and 98.7% of calls resulted in a patient being transported to hospital. 50% of calls for chest pain were made within 60 minutes of symptom onset and the median time between symptom onset and arrival at hospital was 1 hour 45 minutes.

Conclusions

In our jurisdiction, a large number of patients call an ambulance for chest pain, and the epidemiology appears similar to that described for other populations. A large proportion of chest pain patients call triple zero within a short time-frame, and therefore may gain great benefit from very early reperfusion therapies such as prehospital fibrinolysis. This collaborative research has the potential to inform strategic health priorities.

THE IMPEDANCE THRESHOLD DEVICE IN ADULT CARDIAC ARREST PATIENTS

PM Middleton, SR Davies

Ambulance Research Institute, Ambulance Service of New South Wales

Background

The impedance threshold device (ITD) is a valve used in cardiopulmonary resuscitation, which limits air entry into the lungs during chest wall recoil between compressions, and is designed to reduce intra-thoracic pressure, enhancing venous return to the heart, cardiac output and cerebral blood flow.

Objectives

To evaluate the current evidence for the use of an ITD as an adjunct to CPR in the resuscitation of adult cardiac arrest patients.

Methods

A systematic and highly sensitive search for relevant published literature was conducted using a combination of database-specific keywords and text words in Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials, Medline and Embase. As an adjunct to electronic searches, reference lists of all relevant studies were examined, and forward and backward searching of relevant papers was performed in Scopus. An internet-based search for grey literature was undertaken. Eighteen studies met the criteria for inclusion in this review.

Findings

Evidence of improvement in long term survival rates has not been demonstrated, however when the ITD is used as an adjunct to CPR in adult cardiac arrest patients an improvement in haemodynamic parameters and rates of return of spontaneous circulation has been reported. Limitations in study design, combined with a paucity of studies in cardiac arrest patients mean that these improvements in short term outcomes cannot be confidently generalised to the broader cardiac arrest population. One large, well-designed, randomised controlled trial of the ITD in out-of-hospital cardiac arrest patients was terminated early due to a lack of evidence of intervention effect.

Conclusion

This systematic review was unable to find evidence of any beneficial effect associated with the use of an ITD in adult cardiac arrest patients.

SUPRAGLOTTIC DEVICES AS AN ALTERNATIVE TO BAG VALVE MASK VENTILATION IN CARDIAC ARREST PATIENTS

Paul Middleton, Suzanne Davies

Ambulance Research Institute, Ambulance Service of New South Wales

Background

Endotracheal intubation is considered the optimal method for securing the airway during cardiovascular resuscitation; however maintenance of intubation skills is difficult without regular experience. As a result, bag-valve-mask ventilation has become the default technique for initial airway management; however it also requires skill and is physically demanding when performed effectively. Concerns have also been raised about the potential for gastric insufflation and regurgitation. Supraglottic airways offer the potential for safe, effective, single-operator airway management that may be used by less experienced operators.

Objectives

To evaluate the current evidence comparing supraglottic airways, to bag-valve-mask ventilation alone, for airway management in adult cardiac arrest patients.

Methods

A systematic search for relevant published literature was conducted using a combination of database-specific keywords and text words in The Cochrane Library, Medline and Embase. Reference lists of all relevant studies were examined, and forward and backward searching of relevant papers was performed in Scopus. An internet-based search for grey literature was undertaken. Twenty-nine studies met the final criteria for inclusion in this review.

Findings

There is no high-level evidence that the use of a supraglottic airway device, compared to a bag-valve-mask, improves any outcomes for adults in cardiac arrest. Studies performed in mannequins or fasted, anaesthetised patients, support the use of a supraglottic airway for improved ventilation and decreased gastric insufflation; however these cannot be considered valid models of cardiac arrest. There is a trend for improved ventilatory outcomes with newer supraglottic devices. No adverse outcomes have been reported with the use of supraglottic devices.

Conclusion

Supraglottic airways may offer a safe and effective alternative to bag-valve-mask ventilation; however the absence of any well-designed studies comparing these airway management strategies in the cardiac arrest population precludes the recommendation of either strategy.

PRE-HOSPITAL EPIDEMIOLOGY OF SUSPECTED STROKE PATIENTS IN NEW SOUTH WALES, AUSTRALIA

Paul Middleton¹ R Markus² S Anand¹ M Longworth² J Simpson³ M-L McLaws⁴ J Patterson¹

¹Ambulance Research Institute, Ambulance Service of New South Wales

²Greater Metropolitan Clinical Taskforce Stroke Network

³Sydney School of Public Health, University of Sydney

⁴School of Public Health and Community Medicine, University of New South Wales

Introduction

Effective treatment of patients with stroke requires a coordinated effort between prehospital and in-hospital care. As effective t-PA therapy depends on administration within 3 hours of the onset of symptoms of stroke, patient outcomes largely depend on prompt recognition of stroke and shorter transport times by ambulance services. This paper describes the prehospital epidemiology of stroke in NSW, Australia.

Methods

This retrospective study utilised linked data from CAD (Computer Aided Dispatch) and PHCR (Patient Health Care Record) databases from the Ambulance Service of NSW. Data for the time interval July 2004 to June 2007 were studied. Descriptive statistics were investigated using SPSS. Spatial description of suspected stroke patients was carried out using MapInfo.

Results

32,304 emergency calls were received and classified as potential strokes for the time interval June 2004 and July 2007 by Emergency Medical Dispatchers (EMDs), paramedics or both. There were 17,669 (54.7%) cases categorised as priority 1 & 14,635 (45.3%) cases categorised as priority 2, and patients included were aged between 18 & 111. In Metro NSW there were 12,293 (54.4%) Priority 1 calls & 10,313 (45.6%) Priority 2 calls, and in Regional NSW there were 5,330 (55.3%) Priority 1 calls & 4,300 (44.7%) Priority 2 calls. 78.9 % of metro patients were transferred to Stroke Care Unit. Agreement between EMDs and paramedics was found to be 42.3%. 13,531 (83%) of metropolitan patients were taken to a Stroke Care Unit. We calculated that a patient has to be within a 43 kilometre radius of a Stroke Care Unit to receive t-Pa within 3 hours.

Conclusion

Approximately 10,000 patients each year are attended with a presumptive diagnosis of stroke, of which two thirds are in metropolitan areas and one third in rural areas. More than half of cases in both distributions are treated as a high priority, and the majority of patients are taken to hospitals with stroke care units where they are available, and we have calculated an effective maximum distance which will allow effective lytic treatment.

UTILITY OF GEOSPACIAL INFORMATION AND MAPPING TO BETTER UNDERSTAND AMBULANCE OPERATIONS

Paul Middleton, Jillian Patterson, Sowmya Anand, Jason Bendall

Ambulance Research Institute, Ambulance Service of New South Wales

Background

Ambulance jurisdictions collect large quantities of data that at times can be difficult to visually display in a readily interpretable format. Recent advances in geospatial mapping help visualise emergency medical service (EMS) epidemiological data in meaningful ways. Mapping of EMS incidents adds the dimension of geographic analysis to the data which makes it easy to present important information to key decision-makers quickly, efficiently and effectively.

Objectives

The aim of this study was to apply mapping techniques across a range of ambulance data sets in order to describe geospatial trends in ambulance data.

Methods

Geospatial information routinely recorded from ambulance vehicles was analysed using SAS 9.1 and MapInfo Professional 10.0, to obtain counts, rates and proximity to resources. Descriptive statistics of disease incidence within several kilometers of resource availability were also analysed. Bayesian smoothing (a technique used to minimise the effect of such small changes in counts by using information available from surrounding locations) was used to allow for local variations in rates in areas with small populations such as rural and remote locations.

Findings

Mapping EMS incidents helps to visualise geographic distribution of ‘disease’, identify populations at risk, identify risk factors, analyse trends and may help in the allocation of ambulance resources. Mapping is a powerful tool to enable high level statistical data to be presented in a readily accessible format. This methodology has been applied to conditions including falls, acute coronary syndromes, cerebrovascular disease and vehicle accidents using data from the Ambulance Service of NSW.

Conclusions

While still a relatively new area in pre-hospital research, geospatial mapping techniques shows great promise as a tool for understanding ambulance workload, evaluating resource allocation and is very likely to assist in future service planning efforts.

COMMUNITY PERCEPTIONS OF THE PROFESSIONAL STATUS OF THE PARAMEDIC DISCIPLINE

Pauline Murcott, Brett Williams, Ameer Morgans, Malcolm Boyle

Department of Community Emergency Health and Paramedic Practice, Monash University,
Melbourne

Background

Since its inception, the evolution of prehospital emergency care has seen the traditional role of a paramedic shift from an ‘ambulance driver’ to an emergency healthcare provider who is an autonomous clinical problem solver. Despite proactive steps towards professionalism, the prehospital emergency care sector has not gained ‘professional’ status among other healthcare professions and government bodies.

Objectives

To examine community perceptions of the ‘professional’ status of the paramedic discipline and explore which attributes community members perceive as being most desirable for paramedics.

Methods

Two structured focus groups were conducted with volunteer community members (n=10) from regional and metropolitan Victoria. Participants were asked a range of questions, including what they ‘called’ paramedics, whether they considered the paramedic discipline a profession, and what they perceived as desirable paramedic attributes. Thematic analysis and incidence density qualitative data analysis methods were used to examine the responses.

Findings

The results suggest that the paramedic discipline is perceived as a ‘profession’ by the general community. Participants consistently reported the following attributes as most desirable for paramedics: high levels of skills, caring, empathetic, non-judgemental, culturally sensitive, quick thinkers, inclusive (including family members) trustworthy, honesty, good communicators, highly educated, maintain confidentiality, physical fitness and being well groomed. Interestingly, community members used a variety of titles in referring to paramedics, these included: ‘ambulance drivers’, ‘ambulance officers’, ‘ambos’, and ‘paramedics’, although it was clear that the term paramedic confused many participants.

Conclusions

The general community recognises the paramedic discipline as a profession. Interestingly, the term paramedic is still not synonymous among community members, suggesting further lobbying and role clarification is required. The public perceptions of the paramedic role and the desirable attributes of a paramedic provide important information and an extra dimension regarding curriculum development, core competencies, and establishment of standardised curriculum. Further examination of community perceptions is required, particularly as the discipline struggles with national standards, registration and accreditation.

MENTAL HEALTH CRISES IN THE PREHOSPITAL SECTOR

Pauline Murcott, BAppSc (Human Movement) MBusiness (Event Management)
Jade Sheen, BA.BAppSc (Psych) (Hons), MClinFamTh, DipPsych (Health) **Kathryn Eastwood**, BSc, RN, DipAmbParamedStudies, BParamedStudies, MEmergHlth (MICA), GradDipEmergHlth (MICA), **Leanne Boyd**, BNursing, GCert (Critical Care), PhD **Matt Johnson**, DipAmbParamedStudies, GradDipEmergHlth (MICA) **Kevin De Costa** RN

Department of Community Emergency Health and Paramedic Practice, Monash University, Melbourne

Background

Definitions of a mental health crisis focus on 'risk to self and others'. In the prehospital environment, Ambulance Paramedics are often first on scene and are required to provide assessment and management, in addition to Police and Crisis Assessment Teams. Unfortunately, the management of mentally ill patients in the prehospital sector is often fragmented, with poor interagency communication and liaison evident. These issues were the focus of an educational program created by Monash University and provided the impetus for this literature review.

Objectives

- Review research pertaining to the prehospital management of mental health crises.
- Discuss limitations in the management of mental health crises, drawing from the experiences of consumers, carers and clinicians.
- Provide recommendations to promote optimal patient care.

Methods

A literature search was completed using eight computerized databases. 28 articles met the inclusion criteria. Inclusion criteria consisted of treatment and management during a mental health crisis in the prehospital environment or emergency department within Australia. International research was excluded.

Findings

There is a paucity of literature pertaining to the prehospital management of mental health crises. The data available indicates negative consumer experiences related to perceptions of poor staff attitudes, a lack of training and service limitations. These experiences can have long-term impact. The management of mental health crises within Victoria has also been flagged as problematic, with limited interagency liaison, education and interagency training identified, as well as a limited understanding of consumer and carer experiences.

Conclusions

Future research is required identifying optimal prehospital management of mental health crises. Review of paramedic attitudes towards mental illness and the impact this can have on assessment and management is required. Ongoing mental illness education, increased opportunity for interagency learning and clarification regarding interagency protocols will also promote optimal patient care.

IS EVERY CHEST PAIN A CARDIAC EVENT? A LITERATURE REVIEW OF RISK PREDICTION MODELS FOR THE PREHOSPITAL SETTING

Ziad Nehme¹ BEmergHlth(Paramedic)(Hons), (PhD Candidate) **Malcolm Boyle**² ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClInEpi, PhD **Ted Brown**¹ PhD, MSc, MPA, GradCert(HealthProfEd), BSc(OT), OT(C), OTR, AccOT

¹Department of Occupational Therapy, Monash University, Melbourne Australia

²Department of Community Emergency Health and Paramedic Practice, Monash University, Melbourne, Australia

Background

Optimal patient outcomes from acute coronary syndromes (ACS) depend on rapid diagnosis, accurate risk prediction, and effective implementation of prehospital and in-hospital treatment strategies. While cardiac risk prediction is widely used in various clinical settings, its potential role as part of current contemporary management strategies within the prehospital environment is yet to be understood.

Objectives

The objective of this study was to determine the diagnostic accuracy of a prehospital-friendly short-term risk prediction model for patients presenting with suspicion of an ACS.

Methods

A literature review of 11 electronic medical databases was conducted from 1990 to the end of March 2010 for all English-literature studies. Articles were included if they 1) enrolled an unselected cohort of suspected ACS patients aged >18 years; 2) investigated variables including patient history or physical examination, with or without electrocardiography; 3) provided ≤ 30-day outcome measures; 4) were not computer-based and; 5) were not reliant on tests unattainable in the prehospital setting. Study quality and extracted data were assessed independently by the authors using a standardised data extraction form. Reference lists of relevant articles were also reviewed.

Findings

The search identified 47,645 articles. A total of 12 studies met the inclusion criteria encompassing 73,967 patients. Only one study independently validated the prediction model using 2 external validation cohorts with clinically different interventional outcomes. A total of 2 studies reported statistical modelling including performance testing. Sensitivity and specificity values were reported in 3 studies, ranging from 86% to 100%, and 53% to 96% respectively. A combination of physical examination and patient history performed best against other measures.

Conclusion

Emergency medical services have an important role to play in the identification of short-term outcomes in patients with ACS. Only one study was methodologically sound to produce reliable outcomes, however it requires prospective validation in an Australian population.

CAN RURAL PLACEMENTS LURE PARAMEDIC STUDENTS INTO WORKING IN THE BUSH?

Ziad Nehme¹ BEmergHlth (Paramedic) Hons, PhD Candidate **Ruth Hosking**² BNursing, BHlthSc (Public Health)

¹Department of Occupational Therapy, Monash University, Melbourne Victoria

²School of Public Health, Queensland University of Technology, Brisbane Queensland

Background

In light of significant shortages of rural paramedics, some authors have suggested that graduates' intentions regarding their choice to work in a rural environment can be influenced by the exposure they gain as a student.

Objectives

We sought to investigate what effect rural clinical placement or rural education has on the intentions of paramedic students to practice in a rural environment.

Methods

A review of the literature was conducted on an electronic rural and remote health database devised by the Monash University School of Rural Health. All English-literature articles from 2000 to present were searched using a simple search methodology combining the truncated text terms 'student' or 'placement' or 'employ' or 'education'. Articles met the inclusion criteria if they investigated the value of rural clinical placements or rural education in influencing employment decisions within an Australian context. Articles were not limited to the paramedic discipline, rather collaborated evidence from all health disciplines. Government and peak body publications, as well as reference lists from the articles found where also searched for inclusion criteria.

Findings

A total of 359 articles were located, of which 10 met the inclusion criteria. A further 6 studies were located by searching reference lists. The bulk of the literature related rural health challenges in the area of medicine and nursing, with only one study located that related to the paramedic discipline. Recurring issues related to health student's intent on working in a rural environment are: 1) self-perception of clinical competency; 2) preparedness to practice; 3) diversity of skills mix; 4) incentives for rural living; 5) presence of strong preceptorship models and; 6) rural community exposure.

Conclusion

The decision to work in a rural environment appears to have both personal and professional implications for students. There is a scarcity of data relating to the paramedic discipline.

BAR-CODE SCANNING OF MASS CASUALTIES; TESTING OF AN AUSTRALIAN SOLUTION TO DISASTER PATIENT TRACKING

Dr. Ian Norton¹ Dr. Brendan Morrissey²

¹Director Disaster Preparedness and Response, National Critical Care and Trauma Response Centre ²Emergency Registrar, Royal Darwin Hospital, Australia

Background

During a mass casualty incident (MCI), the new nationally agreed SMART tag triage cards will be used to label patients. Barcodes pre-printed on these tags offer a novel opportunity for early patient number tracking.

Objectives

This study compared normal information flow from scene to ambulance command via radio and face-to-face communication, compared to a parallel functioning, but blinded next-G enabled barcode scan system, reporting to a specifically designed website. We aimed to prove ease of use, with clear and rapid transfer of scene information to the health commanders on site and in the emergency operations centre.

Methods

We used barcode scanners in specially designed PDAs to electronically track patients at a simulated MCI. First responders were shadowed as they triaged patients, the triage tags being electronically scanned and the data collected transmitted to a protected website, available to adjudicators in both the Incident Command Centre and a remote Emergency Response Centre. The accuracy of the information collected by the PDAs was then compared to that reported by first responders, who had no knowledge of the scan results. Patients were rescanned during any change in triage status, or any movement within the disaster area, from incident site to Casualty clearing post to transport loading and to final destination. All scans were instantly sent via an inbuilt sim card to a purpose built website, and recorded all individual triage categories in each location, with updates every 5 seconds.

Findings

71 patients were tagged during the simulation. Information on patient numbers was available electronically 86 seconds after the arrival of first responders on scene, compared to 17 minutes by conventional methods. A total of 54 patients and 17 dead were scanned, whilst a total of 52 patients were reported to the incident command centre with no information available on numbers dead. The average scan time was 6 seconds. The error rate in scanning was 0.84%.

Conclusion

This is the first trial using bar-code scanning technology on SMART cards in Australia. The results indicate that the electronic tagging of patients improves upon the accuracy and availability of information to the emergency services as compared to conventional methods. The electronic system was less prone to human error and provided identical information to the command site and at remote locations. The PDA software proved to be intuitive and simple to use. Phase two trials will be used in larger scenarios, in remote locations using satellite technology, and in other mass gathering events and responses.

TRANSPORT NOT REQUIRED: A LITERATURE REVIEW OF PREHOSPITAL MANAGEMENT OF HYPOGLYCAEMIC PATIENTS

Alexander Olaussen, Tom Banks, Brett Williams

Monash University, Department of Community Emergency Health and Paramedic Practice,
Melbourne, Australia

Background

Hypoglycaemia is a medical emergency that is commonly encountered by paramedics. The definitive treatment for restoring euglycaemic values, irrespective of conscious level, is often facilitated by paramedics. Therefore, the debate regarding leaving the treated hypoglycaemic patient at home has emerged. Current evidence is diverse and current paramedic clinical guidelines are ambiguous. Therefore a literature review has been conducted. Expectantly this work will result in recommendations as to how paramedics can deviate from traditional practice and move towards a new paradigm: *'treat, educate and refer'*.

Objectives

The objective of this study was to review the literature regarding prehospital appropriateness of safely not transporting hypoglycemic patients after treatment.

Methods

Selected electronic databases plus a hand search for grey-literature was conducted ranging from their commencement to the end of May 2010. Inclusion criteria were any study reporting hypoglycaemia, treatment, and transport in the prehospital setting. References of relevant articles were also reviewed.

Findings

The search located approximately 1300 articles, 22 articles met the inclusion criteria. The majority of articles found that it is safe to *'treat and release'* hypoglycaemia in the known diabetic. A high proportion of studies found that this practice is preferred by most patients and reduces the burden on emergency departments. Two studies found that the degree of adverse secondary events was equal in the transported and non-transported groups. Criteria for safely not transporting have been suggested throughout the literature, all of which aim to ensure that every patient requiring transport is transported (high sensitivity). However, if a screening tool is not critically applied, too many patients may be transported (low specificity).

Conclusion

Whilst the management of hypoglycaemic patients in the prehospital setting in Australia appears sufficient, no national study confirms this. Moreover this practice is based more on experience rather than on empirical evidence. Further research is required to examine whether a paramedic screening tool might offer clinically significant differences.

INTRAOSSIOUS ACCESS IN THE PREHOSPITAL SETTING: A LITERATURE REVIEW

Alexander Olausen, Brett Williams

Department of Community Emergency Health and Paramedic Practice, Monash University

Background

Intraosseous access offers paramedics with an alternative to difficult vascular access sites in critically ill patients, thus potentially saving time in gaining access and administration of medications. Although the majority of Australian paramedics use the Manual Intraosseous Infusion Technique (MAN-IO), several other semi-automatic devices are now available, such as Bone Injection Gun (BIG) and Semi-Automatic Intraosseous Infusion System (EZ-IO). Given the choice of devices now available questions have been raised regarding success rates, accuracy, decay of skills, and adverse events.

Objectives

The objective of this study was to review the literature regarding intraosseous insertion/infusion in the prehospital setting.

Methods

Selected electronic databases plus a hand search for grey-literature was conducted ranging from their commencement to the end of May 2010. Inclusion criteria were any study reporting intraosseous insertion and/or infusion (adult and paediatric) by paramedics in the prehospital setting. References of relevant articles were also reviewed.

Findings

The search located approximately 2100 articles, 30 articles met the inclusion criteria. The majority of articles found that intraosseous access (regardless of technique) offers a safe and simple method for gaining access to the patient's circulation. A high proportion of studies found semi-automatic devices offered better and faster intraosseous access compared to manual devices while also producing fewer complications. Findings suggest semi-automatic devices can reduce insertion times and number of insertion attempts when contrasted with manual insertion techniques. Results also suggest these devices are user-friendly, undemanding to teach and easy learn.

Conclusion

Whilst manual intraosseous techniques are currently used by the majority of Australian paramedics, available evidence suggests that semi-automatic devices may provide effective and quicker intraosseous access. Further research is required to examine the level of paramedic skill decay and whether comparative devices offer clinically significant differences.

THE INFLUENCE OF RESEARCH ON THE POLICY AND PRACTICE OF AMBULANCE SERVICES IN THE UNITED KINGDOM: LESSON FOR AMBULANCE AUTHORITIES AND PARAMEDICS

Peter O'Meara,¹ Malcolm Woollard²

¹Charles Sturt University, School of Biomedical Sciences, Bathurst, Australia ²Coventry University, Faculty of Health and Life Sciences, United Kingdom

Background

Ambulance services recognise the importance and value of using evidence in their decision-making. This shift has challenges given the paucity of robust evidence to support pre-hospital interventions and the dearth of strong skills to interpret research findings. This study examined how and to what extent policy makers in United Kingdom Ambulance Trusts use evidence to inform their decision-making.

Objectives

1. Determine the extent to which UK ambulance services have changed their service delivery models, organisational structures and cultures in response to pressures to become evidence-based providers of healthcare.
2. Understand how higher education institutions have influenced changes to ambulance delivery models, organisational structures and cultures in the UK.

Methods

Data was collected and analysed using a 'process model' of evidence utilisation to examine the translation of research into policy and practice. Semi-structured interviews were conducted with established academic researchers in paramedic practice / ambulance systems, members of the College of Paramedics, and key decision makers within each of the participating Trusts.

Findings

Decision making in UK Ambulance Trusts is partly driven by evidence. Researchers see varying levels of research engagement and turning copious data into good research is a challenge. Trusts have appointed research leads and formed a collaborative National Ambulance Research Steering Committee. However, there remain few research educated paramedics and managers who are ready to embrace evidence based practice.

Conclusions

One of the major challenges is how ambulance services and researchers manage their relationships when they may have very different views of the world, with many researchers placing 'evidence' above other considerations such as public confidence and organisational image that managers are required to consider when making policy decisions. This clash of values revolves around their respective concepts of what evidence is and the importance they attach to the influence of context in the decision-making processes.

INCIDENTS OCCURING ON WESTERN AUSTRALIAN BEACHES

Chris Peck¹, David Reid²

¹Operations Manager, Surf Life Saving Western Australia ²Westpac Life Saver Helicopter Rescue Service, Surf Life Saving Western Australia

Background

Surf Life Saving Australia (SLSA) is Australia's premier coastal surveillance and rescue service. For more than 100 years lifesavers have patrolled Australian beaches, saving lives. In 2008-09 over 40 000 Australian surf life savers and lifeguards conducted more than 25 000 rescues and treated over 69 000 people for injuries and illnesses.

Objectives

The objectives of this study were to:

- analyse the trends in reported incidents occurring on patrolled Western Australian beaches, and review what proportion of those injured were transported to hospital by ambulance; and
- study underlying factors present at the time of the reported incidents.

Methods

Non-identifiable data generated from SLSA Incident Report Forms and input into the Incident Report Database was extracted. Microsoft Excel was used to summarise and analyse the information produced.

Findings

Over the four financial years 2006-07 to 2009-10 there were 1301 reported incidents, an increase of 63% when reviewed on an annual basis. The majority of incidents were public first aid (999, 77%). 175 persons were transported to hospital by ambulance. The majority of ambulance transports were for suspected spinal injuries (35%), dislocations (14%) or suspected fractures (12%).

Males (57%) were more likely than females to be involved in incidents. Children (aged 0 to 14) made up one-third of those involved in incidents, and 22% of those transported to hospital by ambulance. Of the incidents reported, the majority occurred outside of, but near the flagged area (42%). Incidents were most likely to occur when the weather was fine (84%).

Conclusions

There has been an upward trend in the number of incidents attended by surf lifesavers. Of the persons assisted, 175 were transported to hospital by ambulance, the majority with suspected spinal injuries.

1. 2009-10 Coastal Drowning Report.
http://www.beachsafe.org.au/Low_level_info/Drowning_Reports

INJURIES AND ILLNESSES OCCURRING ON WESTERN AUSTRALIAN BEACHES

Chris Peck¹ David Reid²

¹Operations Manager, Surf Life Saving Western Australia ²Westpac Life Saver Helicopter Rescue Service, Surf Life Saving Western Australia

Background

Surf Life Saving Australia (SLSA) is Australia's premier coastal surveillance and rescue service. For more than 100 years lifesavers have patrolled Australian beaches, saving lives. In 2008-09 over 40 000 Australian surf life savers and lifeguards conducted more than 25 000 rescues and treated over 69 000 people for injuries and illnesses.¹

Objectives

The objectives of this study were to:

- analyse the types of injuries and illnesses occurring on Western Australian beaches; and
- examine the types of injuries and illnesses which required referral to medical care

Methods

Non-identifiable data generated from SLSA Incident Report Forms and input into the Incident Report Database was extracted. Microsoft Excel was used to summarise and analyse the information produced.

Findings

Over the four financial years 2006-07 to 2009-10 there were 1301 reported incidents on Western Australian beaches.

Of the 1100 incidents which had an injury or illness recorded, 80% were traumatic in nature. 29% of injuries and illnesses were open wounds and 13% were marine stings. 4% were respiratory problems, and 1% cardiac problems.

Of the 1091 incidents where a referral destination was recorded, 49% of persons were not referred to further care. 19% were referred to a medical practitioner and 13% to hospital by their own transport. 16% of persons were transported by ambulance.

Over the four year period, 30% of persons referred to medical practitioners had open wounds, the same proportion of persons referred to hospital by their own transport. 175 persons were transported to hospital by ambulance. Of the persons transported by ambulance, 35% had suspected spinal injuries, 14% dislocations, and 12% suspected fractures.

Conclusions

The majority of medical incidents attended by surf lifesavers are traumatic in nature. Around half the persons treated are not referred for further care, with around a third referred for further care by their own transport.

1. 2009-10 Coastal Drowning Report.

http://www.beachsafe.org.au/Low_level_info/Drowning_Reports

THE DEVELOPMENT OF A REVISED PREHOSPITAL SEARCH FILTER FOR THE COCHRANE LIBRARY

Sarah Piper BExercise and Sport Science **Stephen Burgess** BHthSc, GradDip EmergHlth (MICA), MPH, PhD Candidate **Erin Smith** BHlthInfoManagement, MCLin. Epi, PhD

Department of Community Emergency Health and Paramedic Practice, Monash University, Melbourne, Australia

Background

Prehospital care literature continues to increase as the field evolves, reflecting changes in education and clinical practice. Reviews of the prehospital based research identify gaps in the evidence for current practice and policy. There are also issues with the accessibility of the literature. In an attempt to remedy these problems a prehospital search filter for the Cochrane Library was developed and first published in 2004.

Objectives

The objective of this study was to examine the Cochrane Library using an updated and expanded set of search terms to establish whether new search terms should be added to the existing search filter to improve the number of positive results.

Methods

The authors developed and tested a list of new search terms for an updated prehospital search filter for the Cochrane Library. A reviewer searched the Cochrane Library 2010, issue 2 using the updated search strategy terms to identify reports that were not detected by the first version of the prehospital search filter. Three reviewers screened the additional titles and applied predetermined criteria to identify relevant prehospital based reports. These reports were then categorized as randomized or nonrandomized trials.

Findings

The prehospital search filter for the Cochrane Library version 2.0 returned 38,648 reports, an additional 1,604 reports compared to prehospital search filter for the Cochrane Library version 1.0 which returned 37,044 reports. Further analysis found that only 20 were identified as relevant to the prehospital field.

Conclusions

The prehospital search filter for the Cochrane Library version 2.0 has improved sensitivity compared to the prehospital search filter for the Cochrane Library Version 1.0 however, specificity is only marginally improved. Nevertheless, as the prehospital field suffers from a chronic deficiency of relevant, high quality publications, prehospital search filter for the Cochrane Library version 2.0 is a valuable tool for searching the prehospital literature.

HUMAN FACTORS IN PREHOSPITAL CARE – AN UNEXPLORED WILDERNESS

Roger Price, Jason Bendall, Paul Middleton

Ambulance Research Institute, Ambulance Service of New South Wales

Background

An understanding of human factors that contribute to errors, near misses and adverse patient events is an important part of providing good quality health care. ‘Human factors’ research investigates the role of systems, culture, technology and people in behaviour. Risk in prehospital care is unquantified but may represent one of the riskiest areas in health; this paper investigates the role that human factors play in patient safety in prehospital care.

Objectives

To understand the nature of the impact of human factors on adverse events in a large Australian ambulance jurisdiction.

Methods

This was a qualitative study investigating serious reported incidents involving the Ambulance Service of NSW between 2005 and 2009. A thematic analysis was undertaken and themes and trends were identified.

Findings

There were 257 serious incidents identified; delays, omission of clinical procedures and communication breakdowns were recurrent themes in severe adverse events. Lack of resources, rural locations, non-conveyance and non-English speaking patients were also implicated as risk components. Further work is now under way to identify fundamental issues and causes underlying these risk factors, using an online survey and further incident data mining.

Conclusion

Paramedics work within a complex health system, in uncontrolled environments, under time pressure and routinely have to deal with interruptions, uncertainty and distractions. Key themes in adverse incidents were found to be delay, omission of clinical procedures and breakdown in communications. A human factors approach and a safety culture within the Ambulance sector will help to prevent, mitigate and protect against the synergistic effects of these risk components.

UTILISATION OF AMBULANCE TRANSPORT IN HIGH ACUITY RURAL PATIENTS - MEASURING UNMET NEED

Buck Reed¹ Jason Bendall² Jillian Patterson²

¹Ambulance Service of NSW² Ambulance Research Institute, Ambulance Service of NSW

Background

The use of ambulances has been measured thoroughly by ambulance authorities as part of normal operational data collection. Anecdotally, ambulance usage in rural and remote locations is considered to be lower than in metropolitan areas. However, little data has been collected in terms of the number of ambulance non-users especially in rural and remote areas.

Objectives

The objective of this study was to determine the number of patients accessing rural and remote hospitals emergency departments who were not transported by ambulance and determine the potential level of unmet need in acute patients.

Methods

Hospital emergency department presentation data was obtained for a 12 month period from Hunter New England Area Health Service. This data was analysed for a range of factors including patient triage category and method of arrival to the ED. It was established the patients triaged with an Australian Triage Score of 1 or 2 would likely benefit from ambulance intervention prior to arrival at hospital. A number of individual presentation types with good evidence for improved outcomes from prehospital intervention were also analysed. The location of hospitals was stratified using the ARIA scale to determine the rurality of the facility and population.

Findings

Rural and remote hospitals were significantly more likely to have patients with high levels of acuity present by methods other than ambulance (normally private vehicle) than metropolitan hospitals. Triage category 1 and 2 patients presenting to rural hospitals were 27.9% less like to use an ambulance in an acute medical emergency than patients in metropolitan emergency departments.

Conclusions

Rurality appears to be a factor in the method by which patients access emergency departments. Further investigation is warranted to determine the rationale for differences in methods of arrival at emergency departments between rural and metropolitan populations.

MOVING FROM A PAPER-BASED TO A WEB-BASED CLINICAL PLACEMENT PROCESS

Alison Reeves¹ Malcolm Boyle¹ ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MClinEpi, PhD
Nathan Stam¹ BSc(Hons), DAPS

¹Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

The organising of clinical placements for undergraduate paramedic students has previously been undertaken manually, with all student preferences, student requirements, such as police and working with children check and vaccinations records, being kept in a primitive quasi database using a spreadsheet. Likewise, placement locations and available shifts were also managed manually with a spreadsheet. Communication with students was managed from the clinical placement co-ordinator's email client.

Objectives

The objective of this paper is to describe the web-based clinical placement program and how it improves clinical placement management.

Methods

The web-based program was created on a secure web platform with a database back-end and web front end. The program is password protected and available through any web browser on any operating system.

Findings

The web-based clinical placement program provides a repository for basic student details. The program is also a repository for all placement location data, this includes hospitals, ambulance services, GP clinics and other placement locations, including available shifts. The program has an inbuilt email system which can send emails to individual students or groups of students depending on the requirement. Students who have not presented all their documentation to the admin staff prior to their placement, like the police check, will get a personalised reminder email. Students get their final placement allocation, in a pdf file, emailed to them. Students are given a timeframe in which to submit their preferences based on the available locations, failure to submit a request means the student gets what locations are left after the allocation of locations to the respondents. Student preferences are taken into consideration by the program during the allocation process with each student getting at least one of their first preferences.

Conclusions

The web-based clinical placement program has simplified the placement process and significantly decreased the time and errors attributed to a manual system.

UNDERSTANDING THE IMPACT OF VEHICLE COLLISIONS USING AMBULANCE GEOSPATIAL AND CLINICAL INFORMATION

Paul Riley¹ Sowmya Anand² Paul Middleton² Jillian Patterson² Jason Bendall²

¹NSW Institute of Public Works ²Ambulance Research Institute, Ambulance Service of New South Wales

Background

Road crashes are estimated to cost the NSW community in excess of \$4,000 million annually. Currently NSW crash data does not utilise data from the State ambulance service, and the utility of emergency medical system data to describe the incidence, specific locations and outcomes of collisions is yet to be determined.

Objectives

To describe the location and severity distribution of vehicle collisions in NSW using geospatial, operational and clinical information from the Ambulance Service of NSW.

Methods

Vehicle collisions were identified from Computer Aided Dispatch (CAD) and Patient Health Care Record (PHCR) databases for the period between 1 July 2008 and 30 June 2009. The search was refined to include only incidents where a patient was conveyed to an emergency department (ED). Injury severity was categorised into one of four categories based on clinical and operational criteria. Data was analysed using MapInfo Professional 10.0 and SPSS 17.0.

Findings

During the study period 9810 incidents meeting the inclusion criteria were identified, with the frequency of estimated severity being 2.3%, 5.6%, 8.7% and 83.4% for serious injury, potentially serious injury, potential injury and other respectively. Maps of incidents by severity classification were generated.

Conclusions

Ambulance jurisdictions are custodians of geospatial and clinical information that can classify injury severity, and identify locations of collisions, that are likely to result in potential or serious injury. Systems that assume that all patients conveyed to an ED are 'serious' potentially over-represent the injury burden associated with vehicle collisions. The availability of these data is expected to assist with prioritisation of interventions for identified 'at risk' locations. This collaborative research has the potential to inform both strategic health and infrastructure priorities.

THE IDENTIFICATION, ASSESSMENT, AND MANAGEMENT OF PSYCHIATRIC PRESENTATIONS BY PARAMEDICS WITHIN THE COMMUNITY

Louise Roberts

Research Higher Degree Candidate, School of Nursing and Midwifery, Faculty of Health Sciences, Flinders University, Adelaide, South Australia.

Background

“I mean I guess we’re taking them to a facility where they’re going to be fully assessed and managed, so my job is to keep them calm, reassured in order to be able to get them to hospital so that they can get the treatment that they need” (second interview: number 10).

The changing role and expectations of paramedics to provide high quality, evidenced-based, and in some cases extended care, coupled with reform in the mental health sector places paramedics in a position that is both complex and challenging. Reform, based on community models of care, advocates increased participation from a wide array of health professionals, including paramedics.

This research asks the question: how do paramedics identify, assess and manage psychiatric presentations in the community. What paramedics do and how they do it is critical to the interaction and care provided to the people they attend. As providers of ‘frontline’ care to the general public paramedics face expectations that they will be able to treat and care for a wide variety of individuals and their needs. The complexity and often long term needs of people with mental health problems frequently challenges this assumed role and personal, organisational and public expectations.

Objectives

The objective of this qualitative research is to produce detailed descriptions and situated accounts of paramedics’ stories, their views, and strategies in relation to their practice when attending psychiatric presentations.

Method

Using ethnographic techniques data was collected through interviews, observation, case card analysis and focus groups. The data was collected over an eleven month period within the emergency department and ‘on the ramp’ of a tertiary public hospital.

Findings and Conclusions

Two case stories are presented to illustrate the major themes of assessment, communication and interaction, education, risk, ideas of safety and caution, and interaction during handover.

IMPROVING MOBILE INTENSIVE CARE FLIGHT PARAMEDIC PRACTICE: DEVELOPING GOAL DIRECTED THERAPY FOR THE INTER-HOSPITAL TRANSFER OF STROKE PATIENTS

Nicholas Roder GradCert (MICA), GradCert (AeroMedical), DipSci (Nursing), Med

Mobile Intensive Care Flight Paramedic, Air Ambulance Victoria

Background

The inter hospital (IHT) transfer of the intubated, ventilated Stroke patient is a common case type for the Mobile Intensive Care Flight Paramedic (MFP) operating within Air Ambulance Victoria (AAV). While Goal Directed Therapy (GDT) has been introduced for the Traumatic Brain Injured and Septic Shock IHTs, Clinical Practice Guidelines (CPG) aimed at standardising the management and physiological targets of Stroke patients have yet to be developed.

Objectives

The objective of the research was to understand the frequency, patient physiological condition and efficacy of MFP management of the ventilated Stroke IHT patient.

Methods

With the consent of AAV, data was extracted from completed Patient Care Records (PCR). The study included patients where the MFP was the clinical leader for hospital to hospital transfers of adult patients (>14 years). The patient's vital signs observed upon the MFP's arrival at the transferring hospital to those recorded at destination were compared. Supporting data was also collected, including (but not limited to) sedation type, infusions, fluid administration and paralysis use.

Findings

Stroke patients represent the 21.1% (n=43) of adult ventilated IHT case type. Upon handover at the transferring hospital, MFPs noted under-sedation in 32.5% (n=14) patients, while 2 required intubation by the MFP prior to retrieval. At destination 79.1% (n=34) of patients had a Systolic Blood Pressure between 120 and 200mmHg, compared to 67.5% (n=29) upon arrival of the MFP at the transferring hospital. At destination 86.0% (n=37) had ETCO₂ readings between 30 to 37mmHg and 97.7% of patients had an SPO₂ >98%. A Noradrenaline or Adrenaline infusion accompanied 11.6% (n=5) of patients, 4 of which were commenced by the MFP.

Conclusions

Retrospective analyses of the frequency, clinical presentations and MFP management of the ventilated Stroke IHT have formed the genesis for the development of a relevant CPG. Furthermore, there is supporting evidence that in the absence of a specific Stroke CPG, MFPs are already actively performing GDT.

IN-HOSPITAL RAPID SEQUENCE INTUBATION BY MOBILE INTENSIVE CARE FLIGHT PARAMEDICS: CAN PARAMEDIC SKILLS BE EFFECTIVELY USED IN-HOSPITAL?

Nicholas Roder GradCert (MICA), GradCert (AeroMedical), DipSci (Nursing), Med

Mobile Intensive Care Flight Paramedic, Air Ambulance Victoria

Background

Approximately 45% of the Mobile Intensive Care Flight Paramedic (MFP) cases within Air Ambulance Victoria (AAV) are dedicated to the inter-hospital transfer (IHT) of critical patients from regional hospitals. Primarily introduced for out-of-hospital incidents, Rapid Sequence Intubation (RSI) has become a component of the MFP's IHT skill set, facilitating safe patient retrieval in the aero-medical environment.

Objectives

The objectives of this study were to understand the frequency, indication for need, success rates, physiological impact and patient case types that required in-hospital RSI by the MFP.

Methods

With the consent of AAV, data was extracted from completed Patient Care Records (PCR). The study included patients where the MFP was the clinical leader for hospital to hospital transfers of adult (>14 years) patients that were intubated for transfer, resulting in a sample of 204 cases over a twelve month period.

Findings

The data revealed that 8.8% (n=18) of the 204 ventilated IHT cases within this study underwent MFP RSI prior to departure from the transferring hospital. The indications for all (100%) the RSIs were supported within the AAV Clinical Practice Guidelines (CPG). Of the 18 cases, 88.9% (n=16) were conducted using pharmacology under AAV's (CPG), while the remaining 2 were inducted using the hospital's protocol. All (100%) of the patients were successfully intubated. Of the 18 patients, 7 had a Glasgow Coma Score ≥ 12 , with conditions ranging from facial burns to Traumatic Brain Injury. The mean at-hospital time for the 18 RSI patients was 51 minutes, which was three minutes greater than the overall mean time (n=204). All patients demonstrated improvements in their vital signs recorded on arrival at the destination hospital compared to the first recorded at the transferring hospital.

Conclusions

Despite the broad range of clinical presentations and unfamiliar work setting, MFPs demonstrated they could successfully perform a complex out-of-hospital skill set within the IHT environment, working within the AAV CPGs.

THE GROWTH OF PARAMEDIC ROLES AND THE URGENT NEED FOR NATIONAL STANDARDS

**James Ross, MBBS MMED MPH GCert ehealth GAICD AFACHSE FAsMA FAFPHM
FAFOEM**

Medical Director - Aspen Medical

The traditional role of paramedics in the Australian healthcare system has been as ambulance officers. This role is developing further, with the gradual move towards 'professionalisation' of paramedics. However there is still a dramatic variation in training, skills, definitions and capability of paramedics across Australia.

There is a greatly expanding niche for paramedics in Australia based around remote, often solo, practice. This development introduces new challenges for preparation and management of paramedics, for training, credentialing, scope of practice, clinical oversight, skills maintenance and career development. At present there is no regulatory control over paramedic employment other than that provided by an employer. This places much more responsibility on employers (many of whom do not understand the risks), but also opens up the risk of inappropriate use of paramedics without sufficient support structures and/or outside of a defined scope of practice. Many employers are increasingly employing paramedics in such roles.

Having scoped the issues facing the paramedic profession, this presentation discusses one program being implemented to provide clinical governance covering the use of 'remote paramedics', including plans for use of paramedics both domestically and internationally.

PARAMEDICS, END OF LIFE DECISION MAKING AND THE LAW: GOLD STANDARD CARE THROUGH COMPLIANCE WITH THE LAW

Brian Sengstock

Doctor of Philosophy, Law and Justice Research Centre, Faculty of Law, Queensland University of Technology, Brisbane, Australia, and Adjunct Senior Lecturer - Paramedic Science, University of the Sunshine Coast, Maroochydore, Queensland, Australia

Background

Paramedics are routinely required to make end of life decisions in the pre-hospital emergency care setting, often with limited knowledge of the patient's preceding medical history or express wishes in relation to life-sustaining measures. To date there has been no research involving paramedics and end of life decision making in an Australian context from a legal perspective. The legal effect of advance health directives on the decision to withhold or withdraw life-sustaining measures in an acute emergency in the pre-hospital care setting is as yet unexplored.

Objectives

The objective of this study is to establish the extent to which paramedics are currently compliant with the law in relation to end of life decision making and the degree to which paramedics understand the operation and validity of an advance health directive in the emergency pre-hospital care setting.

Methods

Doctrinal analysis of legislation and common law will be used to determine the current legal basis for the operation of advance health directives and the legal issues surrounding this. Qualitative data is being collected from paramedic students and paramedics using scenarios in focus group and individual interviews to determine current compliance with the law. This data will be analysed using grounded theory methods.

Findings

Anecdotal evidence to date indicates that paramedics and paramedic students have a limited understanding of the operation and validity of advanced health directives.

Conclusions

The outcome of a decision to withhold or withdraw life-sustaining measures is irreversible and could have legal implications for paramedics and family members involved in the decision to withhold or withdraw life-sustaining measures in the event that the decision is found to be legally unsupported.

ATTITUDES AND PERCEPTIONS OF PARAMEDICS TOWARDS PREHOSPITAL RESEARCH AND EVIDENCE-BASED PRACTICE

Paul Simpson, Jason Bendall, Mark Goodger, Monica Sitkowski, Suzanne Davies, Paul Middleton

Ambulance Research Institute, Ambulance Service of New South Wales

Background

There is rapidly expanding body of research which investigates prehospital care provided by paramedics. Evidence-based practice (EBP) is becoming increasingly common within emergency medical service (EMS) jurisdictions, yet little is known about how paramedics feel about the emergence of this paradigm. As operational paramedics will be asked to change their clinical practice based on prehospital evidence and to participate in research initiatives, a better understanding of their attitudes and perceptions will assist in successful translation of research into practice, and will optimise engagement in research projects.

Objective

To investigate the attitudes and perceptions of paramedics towards prehospital research and evidence-based practice.

Methods

An anonymous, voluntary on-line electronic survey was conducted, in which paramedics were asked to respond to a series of questions relating to prehospital research generally and also to how they thought evidence-based practice and a research-based culture would affect them.

Findings

At the conclusion of the survey period 929 paramedics had responded, representing 25% response rate. High levels of confidence in research influencing clinical practice were reported by 79.9% of respondents and 83.2% believed that the influence would be a positive one. When asked about the likelihood of change in their individual practice based on research results, 74.4% responded that this was at least a likely outcome. When asked about their supportiveness towards research in their ambulance service, 89.6% said they were at least likely to volunteer to participate in the implementation of research initiatives and 73.2% stated they were at least supportive of the development of an evidence based culture.

Conclusions

The majority of paramedic respondents in this population appear to have positive attitudes towards prehospital research and the emergence of evidence-based practice. There appears to be a perception that prehospital research will have a positive influence on clinical practice and will improve patient care. In this jurisdiction, paramedics appear likely to support research initiatives and will volunteer to participate in implementation.

TRICYCLIC ANTIDEPRESSANT OVERDOSE: A REVIEW OF THE PATHOPHYSIOLOGY, PHARMACOLOGICAL INTERVENTIONS AND PREHOSPITAL MANAGEMENT STRATEGIES

Gavin Smith BParaStudies, GradDipEmergHealth, MEH, (PhD Candidate)

Ambulance Victoria, Melbourne, Australia

Background

Tricyclic antidepressant (TCA) overdose represents a high percentage of those patients presenting in the prehospital setting as overdose. Management of these patients is complex, and presents challenges for paramedics due to the toxic nature of these substances, and the propensity for rapid deterioration within six hours of ingestion. A complete understanding of the mechanisms and complications of TCA overdose is essential in order for paramedics to appropriately manage these patients.

Objectives

The objective of this study was to review the available literature to ascertain the pathophysiology of TCA overdose, and to identify specific tools which may be of use in the prehospital setting.

Methods

A systematic review of the electronic literature was conducted. Six clinical studies examining a variety of pharmacological interventions were identified. A single study related to the prehospital emergency care setting.

Findings

TCA overdose has a mortality rate of 70% within six hours of ingestion. There are three primary areas requiring specific management resulting from TCA overdose: Cardiovascular System, Central Nervous System, and Anticholinergic effects. A range of pharmacologic agents have been investigated for specific use in this setting, yet none have been proven definitive. Sodium Bicarbonate 8.4% continues to represent the most popular intervention. Prolonged resuscitation efforts of up to five hours in the cardiac arrest TCS overdose patient were highlighted as potentially beneficial. A useful mnemonic was also identified for the identification of TCA overdose.

Conclusions

TCA overdose will continue to represent a complex and dynamic patient for paramedics. The use of mnemonic aids in assessment should assist in improved recognition of the time critical nature of these patients, and further study into the use and effectiveness of Sodium Bicarbonate 8.4% is required in order to formulate changes in practice, and to reduce mortality and morbidity in this patient group.

PREHOSPITAL TRICYCLIC ANTIDEPRESSANT OVERDOSE: A RETROSPECTIVE CASE SERIES STUDY OF PREHOSPITAL INTERVENTION BY MICA PARAMEDICS IN METROPOLITAN MELBOURNE

Gavin Smith^{1,2} BParaStudies, GradDipEmergHealth, MEH, (PhD Candidate)

Malcolm Boyle² DipBus (GenAdmin), BInfoTech, MCLinEpi, PhD

¹MICA Paramedic, Ambulance Victoria, Melbourne, Victoria, Australia ²Senior Lecturer, Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

Tricyclic antidepressant (TCA) overdose represents 0.01% of those patients presenting to ambulance paramedics as overdose. However, the likelihood of toxicity leading to subsequent cardiac arrest in this group is approximately 70% within six hours of ingestion. Management of these patients is complex, and presents challenges for paramedics in the prehospital setting.

Objectives

The objective of this study was to identify the effectiveness of MICA Paramedic management of TCA overdose using Sodium Bicarbonate 8.4%. This information may prove useful in formulating a specific TCA overdose Clinical Practice Guideline within Ambulance Victoria (AV).

Methods

A retrospective case series study of Victorian MICA Paramedic management of TCA overdose. Data was obtained from the VACIS database (AV) for the period March 2007 to November 2009. Descriptive statistics were used to analyse the data. Inclusion criteria were those cases listed as overdose, TCA, specific drug overdose within the window. Patients were excluded if there was no way of confirming that TCA were involved in the overdose.

Findings

Of the 23264 cases dispatched as overdose, 264 were identified as TCA overdose or polypharmacy overdose likely to involve TCA. During the review period 68 cases, having demonstrated toxicity, required MICA intervention with Sodium Bicarbonate 8.4%. Improvement was noted in 61.1% of cases where intervention occurred. Most notable effects of treatment were a reduction in QRS width on ECG, improved perfusion, and resolution of arrhythmia. Also of note was the failure of patients (ECG or perfusion status) to respond after two doses of Sodium Bicarbonate 8.4%.

Conclusions

MICA Paramedic intervention in TCA overdose using Sodium Bicarbonate 8.4% has merit. Further study is required to define the nature of impact of current intervention on the condition, and to support an evidence based approach to treatment methodology in the form of a specific Clinical Practice Guideline for the future.

USE OF THE VALSALVA MANOEUVRE TO MANAGE SUPRAVENTRICULAR TACHYCARDIA BY MICA PARAMEDICS IN MELBOURNE: A RETROSPECTIVE CASE STUDY REVIEW OF EFFECTIVENESS

Gavin Smith^{1,2} BParaStudies, GradDipEmergHealth, MEH, (PhD Candidate)
Malcolm Boyle² ADipBus, ADipHSc(AmbOff), MICA Cert, BInfoTech, MClinEpi, PhD

¹Ambulance Victoria, Melbourne, Australia

²Monash University, Department of Community Emergency Health and Paramedic Practice, Melbourne, Australia

Background

Recent studies have identified an evidence based model of Valsalva Manoeuvre (VM) performance. The use of the VM for supraventricular tachycardia (SVT) management in paramedic practice is developing structure as this model is integrated, yet to date there has not been an assessment of effectiveness in reverting SVT in the clinical setting.

Objectives

This study sought to identify the effectiveness of the VM in reverting SVT in the prehospital setting by MICA paramedics in Melbourne, Australia. This study also sought to quantify other factors which may impact upon reversion success.

Methods

A retrospective case review utilising data from the Victorian Ambulance Computerised Information System (VACIS) database. The data period was November 2006 to November 2007. Inclusion criteria were patients presenting with SVT in the prehospital setting, managed by MICA paramedics. Exclusion criteria were rhythms other than SVT identified before or during intervention by paramedics.

Findings

The study examined 156 cases, with patient age ranging from 11-98 years, mean age of 56 years (CI 53.71 to 56.55), and median age of 57. Primary dispatch was “chest pain” (50.2%), and “Heart problems/cardiovascular problems” (23.2%). A single VM attempt in 61.4% of cases, with a maximum of seven attempts was noted. Mean time from onset of SVT to VM attempt was 27.76 minutes (CI 26.22 to 29.31), with no statistically significant difference identified by age, gender, cardiac history, history of SVT, or current medications. The study identified a reversion success of 23.2%.

Conclusions

This study has identified current reversion rates using the VM comparable to the emergency medical setting. The results also suggest reversion from SVT is unaffected by gender, age, cardiac history, history of previous SVT, or cardiac medication. A baseline for further studies of the effectiveness of the evidence-based VM in the prehospital setting is established within this pilot study.

THE EXPLORATION OF PHYSICAL FATIGUE, SLEEP AND PSYCHOLOGICAL FACTORS IN PARAMEDICS: A PILOT STUDY

Sarah Sofianopoulos^{1,2} BSc, BEmrgHlth **Brett Williams¹** MHLthSc, PhD Candidate
Frank Archer¹ MBBS, MPH, OAM **Bruce Thompson²** BAppSci, CRFS, PhD

¹Department of Community Emergency Health and Paramedic Practice, Monash University,

²Department of Allergy, Immunology & Respiratory Medicine, The Alfred Hospital,
Melbourne Australia

Background

Paramedics are a group of individuals who face a unique set of challenges, making up a high calibre health care service. With an ever increasing workload paramedics are currently being faced with issues of fatigue and sleep disturbance, which potentially compromises patient and personal safety, physical and psychological health and well-being.

Objectives

To investigate the impact shift work on various aspects of sleep in paramedics.

Methods

Paramedics were asked to complete a number of self-reporting standardised questionnaires. The Epworth Sleepiness Scale (ESS) (8-items), Berlin Questionnaire (BQ) (10-items), Pittsburgh Sleep Quality Index (PSQI) (11-items) and the Beck Depression Inventory (BDI) (21-items).

Findings

The study recruited 60 participants, the majority male 77% (n=46), > 45 years of age 31% (n=19), and having worked shift work between 5-10 years 35% (n=21). Nine out of ten (92%, n=55) of paramedics reported having experienced fatigue in the last 6 months, with 88% (n=53) believing it had affected their performance at work. The ESS reported 30% (n=18) of people had excessive daytime sleepiness, 10% (n=6) being dangerously sleepy. Statistical significance was observed in the ESS items 'chance of dozing while sitting and talking to someone' ($p<0.05$), and 'whilst stopped in traffic for a few minutes' ($p<0.05$). Almost half (48%, n=29) of paramedics answered yes to having nodded off or fallen asleep whilst driving. The PSQI found 68% (n=41) of participants suffered poor quality sleep, while 21 % (n=13) of respondents were at high risk for sleep apnea (BQ). Depression was found to be mild among 27% (n=16) and moderate among 10% (n=6). The questionnaires utilised demonstrated an adequate to excellent reliability with a Cronbach's α ranging between (0.60-0.97).

Conclusion

Shift work affects health and well-being both physiologically and psychologically, which translates from work into home and family life. Further research is warranted to prevent the issues of patient safety, work-related fatigue and the cumulative effects of shift work in paramedic employees.

PREHOSPITAL ASTHMA MANAGEMENT: WHAT DOES THE EVIDENCE SUGGEST – A REVIEW OF THE LITERATURE

Erin Sorensen, Christian Winship, Chloe Bell, Brett Williams, Malcolm Boyle

Department of Community Emergency Health and Paramedic Practice, Monash University

Background

Asthma is an acute respiratory disorder characterised by bronchospasm as a result of stimulation, particularly of sensitive bronchial smooth muscle. As a result of this stimulation, the bronchial smooth muscles contract and constrict causing difficulty in breathing which leads to acute respiratory distress.

Objectives

The objective of this study was to identify the evidence for the prehospital management of acute asthma.

Methods

A literature review was undertaken of the electronic medical databases Ovid Medline, EMBASE, CINAHL Plus, Cochrane Systematic Review, and Meditext from their beginning until the end of April 2010. Keywords used in the search included; asthma, bronchospasm, bronchoconstriction, EMS, EMT, paramedic, emergency medical service, emergency medical technician, prehospital, and ambulance. The keywords were used individually and in combination. Articles were included if they reported on prehospital management of acute asthma or acute bronchospasm/ bronchoconstriction. Articles not written in English were excluded. Inclusion criteria were any study type reporting the management of acute asthma in the prehospital setting.

Findings

Of the 480 papers located, 12 papers met the inclusion criteria. References of relevant article were also reviewed but no additional articles were identified. Of these 17 articles, 8 demonstrated an improved respiratory condition with nebulised beta-agonist, two concluded adrenaline demonstrated no benefit over a nebulised beta-agonist, and two demonstrated that early steroids decreased hospital admission and length of stay.

Conclusion

Evidence suggests that a nebulised beta-agonist and early intravenous steroids is a safe and effective way of managing asthma in the prehospital setting. However, there is still a need for further research into alternative treatment regimes.

PREHOSPITAL THROMBOLYSIS – FEASIBILITY OF IN THE FIELD ADMINISTRATION BY PARAMEDICS IN RURAL AND REGIONAL NSW

Paul Stewart¹, Clare Pemberton², Lindsay Savage³

¹Cardiac Care Project Manager, Ambulance Service of NSW, Sydney, Australia,

²Cardiac Care Project Officer, Ambulance Service of NSW, Sydney, Australia,

³Cardiac Liaison Officer, Hunter New England Health, Southern, Australia.

Background

Acute myocardial infarction (AMI) without treatment has a high mortality and morbidity rate. The success of cardiac reperfusion strategies is significantly influenced by the time interval between symptom onset and restoration of myocardial blood flow. Reducing this time interval is often difficult to achieve, particularly in a rural setting. This project enables ambulance paramedics to provide a clinically proven definitive treatment by administering thrombolytic and anticoagulation treatment to patients with ST Elevation Myocardial Infarction (STEMI) in the field, significantly reducing the time to reperfusion.

Objectives

This project aimed to establish the feasibility of in field administration of thrombolytic medication by paramedics to patients having a STEMI in the setting of rural and regional NSW.

Methods

130 paramedics located in communities where timely access to health facilities was limited were trained and equipped to acquire and transmit a12 lead ECG. Paramedics attending patients with suspected myocardial ischaemia electronically transmit the ECG to a cardiologist for confirmation of STEMI pattern. Paramedics use a developed protocol to deliver thrombolytic and anticoagulation medication, in the case of confirmed STEMI before transporting the patient to the appropriate hospital.

Findings

- The project established the clinical, logistical and communication foundations for the delivery prehospital thrombolysis by paramedics
- Over 840 patients benefited from having a cardiologist interpret their ECG prior to their arrival at hospital
- 45 patients received thrombolytic treatment prior to their arrival at hospital
- No significant adverse events have been associated with this project
- The median time from paramedic arrival at scene to administration of thrombolytic is 28 minutes
- The youngest patient thrombolysed is 37 years of age with the median age being 63 years.

Conclusions

It is feasible for paramedics, assisted by expert review of ECG, to appropriately administer thrombolytic therapy prior to transfer to hospital.

LEGAL AND ETHICAL EDUCATION IN PARAMEDICINE – 'WHY' AND 'HOW'

Ruth Townsend BN, LLB, LLM, GDLP, GCVET, DipParaSc

Lecturer, College of Law and School of Medicine,
and
The Australian National University, Canberra, Australia

Background

There has been a change in the way in which paramedic educational curricula, and delivery of education has been developed in the past few years. One key area that is now included in most programs is that of law and ethics. This understanding of law and ethics is considered to be an attribute of health care professionals that will protect and promote patients' health and well being.

Objectives

The objective of this study was to undertake a meta analysis of literature in the area to examine the optimal way in which law and ethics should best be taught to achieve the desired outcome of a graduate who practices both ethically and legally.

Methods

Literature was gathered from a number of different databases. Due to the lack of material written exclusively in the paramedic discipline, material from similar disciplines including nursing and medicine were also included.

Findings

There is a distinct lack of literature on the teaching of law and ethics in medical education. Despite the fact that law and ethics is now included in most medical/nursing curricula there is little evidence to support its place in curricula aside from aspirations about graduate attributes and there is little evidence to demonstrate that legal and ethical education within curricula alters student to graduate behaviour. There is little evidence to show that legal and ethical education occurs in clinical placement.

Conclusions

There is a need for further research into the 'whys' and 'hows' of teaching law and ethics to paramedic students.

THE SELF-DIRECTED LEARNING READINESS SCALE (SDLRS) FOR PARAMEDIC EDUCATION: RESULTS FROM FOUR UNIVERSITIES

Brett Williams¹ BAVEd, GradCert (ICP), GradDip (EmergHlth), MHIthSc, (PhD Candidate), **Malcolm Boyle**¹ ADipBus (GenAdmin), BInfoTech, MClinEpi, PhD **Richard Brightwell**² BSc (Hons), PhD, CertEd **Scott Devenish**³ MVEEd, BNur, DipParaSc, RN **Amea Morgans**¹ BA.BAppSc (Hons), PhD, GCHE **Graham Munro**⁴ MHSM, BHSc, CCP

¹Monash University (MU), ²Edith Cowan University (ECU),
³Queensland University of Technology (QUT), ⁴Charles Sturt University (CSU)

Background

In paramedic education, self directed learning (SDL) is considered an important characteristic, since paramedics are expected to continue and update their knowledge and understanding throughout their professional career. This is the first study of its kind involving paramedic undergraduate students in Australia.

Objectives

The objective of this study was to determine paramedic students' attitudes and readiness towards self-directed learning.

Methods

A prospective cross-sectional study using a paper survey, the *Self-directed Learning Readiness Scale (SDLRS)*, with a convenience sample of undergraduate paramedic students studying at four Universities in semester 1, 2010. Student attitudes and readiness to self-directed learning were elicited by the *SDLRS* using a 5-point Likert scale (1=Strongly Disagree and 5=Strongly Agree).

Findings

There were 259 students who participated. Most students were from MU (n=113) and CSU (n=77). Two-thirds (n=169) of students were < 25 years of age with 54% female. The *SDLRS* produced important findings in each of its three subscales. The majority of students enjoyed learning new material (M=4.51, SD=0.54), while enjoying the associated challenges that learning brings (M=4.36, SD=0.67). The item 'I can be trusted to pursue my own learning' produced statistically significant findings between year levels, $F(3, 223) = 4.0, p = 0.008$. Post-hoc comparisons found: 1st years (M=4.12, SD=0.87) were significantly different from 3rd years (M=3.80, SD=0.82) and 4th years (M=3.36, SD=1.15). The item 'I am confident in my ability to search out information' also produced significant findings between universities $F(3, 255) = 3.3, p = 0.02$. Post-hoc comparisons found: QUT (M=4.23, SD=0.71) was significantly different from ECU (M=3.67, SD=0.59), CSU (M=3.73, SD=.88) and MU (M=3.75, SD=.82).

Conclusions

As paramedic-orientated degree programs continue to emerge and develop, establishing SDL needs will assist paramedic educators in diagnosing student learning needs, and assist in shaping contemporary and learning-centred curriculum.

VALIDATING THE READINESS FOR INTERPROFESSIONAL LEARNING SCALE (RIPLS) IN UNDERGRADUATE HEALTH CARE STUDENTS

Brett Williams¹ MHIthSc PhD Candidate **Ted Brown**² PhD, MSc, MPA, BScOT (Hons), Grad Cert (HltProfEd), OT(C), OTR, AccOT **Fiona McCook**³ **Jill French**⁴ **Claire Palermo**⁵ **Lisa McKenna**⁴ **Rebecca Scholes**⁶ **Malcolm Boyle**, ADipBus, ADipHSc(Amb Off), MICA Cert, BInfoTech, MCLinEpi, PhD **Janeane Dart**⁵

¹Department of Community Emergency Health and Paramedic Practice, ² Department of Occupational Therapy, ³ Department for Rural and Indigenous Health, ⁴ School of Nursing and Midwifery, ⁵ Department of Nutrition and Dietetics, ⁶ Department of Physiotherapy.

Background

Collaboration and teamwork are now a major focus of the Australian health care system, where teams aspire to optimise outcomes for patients through effective interprofessional collaboration. Interprofessional learning and practice can facilitate the delivery of safe, patient-centered, cost-effective health care. However evidence shows that undergraduates are not well prepared to work interprofessionally, that most healthcare teams report difficulty in working interprofessionally, and that interprofessional practice is a gold practice standard that is rarely achieved. One standardised scale commonly used to measure interprofessional collaboration is the *Readiness for Interprofessional Learning Scale (RIPLS)*.

Objectives

The objective of this study was to assess the construct validity and reliability of the 19-item *RIPLS* in an Australian undergraduate health care context.

Methods

Principal Component Analysis was applied to the *RIPLS*'s 19-items to identify the number of factors followed by Orthogonal Varimax Rotation. A stepwise approach was used to explore the scale's dimensionality. Data were from undergraduate students enrolled in one of seven health-related courses at Monash University - Nursing, Occupational Therapy, Nutrition and Dietetics, Emergency Health (Paramedic), Nursing/ Emergency Health (Paramedic), Physiotherapy, and Midwifery.

Findings

A total of 418 records were analysed. The majority of participants were female (80.4%) and the majority were young adults (<21 years = 25.1%; 21-25 years = 58.1%). Following the rotated factor matrix, three factor extraction criteria were used: Eigenvalue > 1 rule, Scree Test, and Explained Variance (53.8% - items with loadings >.40). This generated a 4-factor solution. Nine items cross-loaded on two or more factors raising questions surrounding subscale validity. The overall 4-factor solution produced a Cronbach's alpha (0.67).

Conclusion

This study raises concerns over the validity and reliability of the 19-item *RIPLS* in its current form for use with Australian undergraduate health students, suggesting further examination of its psychometric properties are required.

A PILOT STUDY EVALUATING AN INTERPROFESSIONAL EDUCATION WORKSHOP FOR UNDERGRADUATE HEALTH CARE STUDENTS

Brett Williams¹ MHLthSc, PhD Candidate **Fiona McCook²** **Ted Brown³** **Jill French⁴** **Claire Palermo⁵** **Lisa McKenna⁴** **Rebecca Scholes⁶** **Malcolm Boyle¹** **Janeane Dart⁵**

¹Monash University Department of Community Emergency Health and Paramedic Practice,

²Monash University Department for Rural and Indigenous Health, ³ Monash University Department of Occupational Therapy,

⁴Monash University School of Nursing and Midwifery, ⁵Monash University Department of Nutrition and Dietetics, ⁶Monash University Department of Physiotherapy

Background

The need for interprofessional education (IPE) is driven by the belief that if professionals are knowledgeable about the skills and roles of each other, it will enable more efficient practice and ultimately optimal patient/client care. Therefore, one of the major focuses of the Australian healthcare system is to promote effective interprofessional learning (IPL) and collaboration.

Objectives:

The objective of this study was to evaluate the effectiveness of a student IPE workshop.

Method:

This study used a repeated-measures design to identify changes in readiness of IPL after the student IPE workshop which involved students from the following disciplines: paramedics, nursing, midwifery, occupational therapy, physiotherapy, and nutrition and dietetics. The independent variable was time, with evaluation occurring before (Time 1), after the student IPE workshop (Time 2) and 6 months following the workshop (Time 3), using the *Readiness for Interprofessional Learning Scale (RIPLS)*. A one-way repeated measures ANOVA was used to identify any changes over time.

Results:

A total of 22 students participated in the study with most students from Paramedics (n=6) followed by Occupational Therapy (n=5) and Nursing (n=5). The majority of students were < 25 years of age (78%), with 73% female. A statistically significant difference was found in four of the 19 items. The item 'Learning with other students will make me a more effective member of a health care team' showed an increase in mean scores across the three time periods, Wilks Lambda = 0.54, $F(2, 17) = 7.17$, $p=0.006$. The item 'Shared learning will help me clarify the nature of patients' or clients' problems' also showed an increase in mean scores across the three time periods, Wilks Lambda = 0.62, $F(2, 17) = 7.17$, $p<0.05$.

Conclusions:

These findings suggest that the IPE workshop for undergraduate health care students is a promising development in improving collaboration among healthcare disciplines. Further work is required in determining if similar IPE workshops should be incorporated into core curriculum.

IS THE AUSTRALIAN PARAMEDIC DISCIPLINE A PROFESSION? A NATIONAL STUDY

Brett Williams¹ MHIthSc, PhD Candidate **Andrys Onsman**² TTC, PhD, Grad. Dip. Ed (TESOL), MEd, BEd, Dip. Teach, **Ted Brown**³ PhD, MSc, MPA, BScOT (Hons), Grad Cert (HltProfEd), OT(C), OTR, AccOT

¹Department of Community Emergency Health and Paramedic Practice, Monash University, ²Centre for the Advancement of Learning and Teaching, Monash University, ³Department of Occupational Therapy, Monash University.

Background

Over the past century, the Australian paramedic discipline has changed dramatically; moving from its origins of an ambulance driver to its current practitioner role and integral member of the Australian health care system. However, at present the Australian paramedic discipline is not considered a full profession.

Objectives

This paper has two objectives: 1) to examine if the Australian paramedic membership views the discipline as a profession and 2) if the paramedic community wants to be considered a full profession within Australia.

Methods

3800 questionnaires were posted to paramedic ACAP members around Australia. An investigation of professionalisation attitudes were investigated using an 8-item paper-based self-report questionnaire using a 5-point Likert scale (1=Strongly Disagree and 5=Strongly Agree).

Findings

A total of 872 paramedics returned completed questionnaires (23% response rate) of which 74% were males. The majority of participants rated highly that the paramedic discipline would benefit from being recognised as a full profession ($M=4.75$, $SD=.54$). The majority of participants agreed that they believed that national registration would not occur within the next 2 years ($M=2.90$, $SD=1.03$). These results are further reinforced with the moderate mean score of 3.10 ($SD=1.10$) that the discipline already exhibits the traits of a professional body. The item whether the 'paramedic sector depends on Higher Education to enhance its chances of becoming a profession' produced significant findings $p=0.04$. Post-hoc comparisons found participants employed between 1-5 years ($M=4.07$, $SD=.88$), and 11-15 years ($M=4.02$, $SD=.91$) was significantly different from those employed > 30 years ($M=3.35$, $SD=1.49$).

Conclusion

The findings from the study suggest two points in relation to professionalism of the paramedic discipline within Australia. Firstly, the paramedic discipline is presently not a full profession and secondly, the paramedic discipline wants to become recognised as a profession. Other professional factors such as national registration, autonomy, national accreditation programs, and the development of a unique body of knowledge require further investigation.

VALIDATION OF THE PARAMEDIC GRADUATE ATTRIBUTE SCALE (PGAS): A RASCH RATING ANALYSIS

Brett Williams¹ MHIthSc PhD Candidate **Andrys Onsman**² TTC, PhD, GradDipEd (TESOL), MEd, BEd, DipTeach) **Ted Brown**³ PhD, MSc, MPA, BScOT (Hons), Grad Cert (HltProfEd), OT(C), OTR, AccOT)

¹Department of Community Emergency Health and Paramedic Practice, Monash University

²Centre for the Advancement of Learning and Teaching, Monash University ³Department of Occupational Therapy, Monash University.

Background

The Australian paramedic discipline sector has seen a remarkable change in a number of areas including education, training, identity, and clinical practice particularly over the past three decades. Therefore preparing future paramedic graduates for these expected changes requires careful alignment of graduate attributes to course curricula, especially as; exactly what ought to constitute such graduate attributes is still a contentious area amongst practitioners. As a first step towards agreement a *Paramedic Graduate Attribute Scale* (PGAS) was devised.

Objectives

The objective of this study was to assess the psychometric properties of the proposed *Paramedic Graduate Attribute Scale* (PGAS).

Methods

Data generated from a national study (n=874) rated the importance of 34 paramedic graduate attribute items. A Rasch Rating Scale Analysis was performed on the 34 items using WINSTEPS (version 3.64.2). Three objectives designed to provide information about the PGAS's construct validity were examined for the best fit model: i) Rasch Model item fit, ii) unidimensionality, and iii) differential item functioning (DIF).

Findings

The Rasch analysis indicated that the PGAS and its 7 factor structure: (Professional Behaviour, Patient Interaction and Welfare, Personal Behaviour and Attitudes, Paramedic and Society, Commitment to Professional and Health Care Outcomes, Scientific Approach to Patient Care, and Clinical Education and Leadership) were valid and reliable. Examination of goodness of fit statistics, unidimensionality, and DIF produced item misfit and item bias in 4 PGAS items. Each of these items was subsequently removed from the PGAS. Item reliability ranged from 0.88-0.99 and variance explained by measures ranged 42.5% to 66.8%.

Conclusions

The PGAS is a psychometrically sound instrument, offering both the industry and university sectors with the first set of empirically-based paramedic graduate attributes that are measureable, valid, and reliable. Based on extensive consultation with industry, the PGAS provides the discipline with a curriculum blueprint necessary for national standardisation, accreditation, and benchmarking of professional paramedic programs.